

SHIP-1 (Phospho Tyr1021) Rabbit pAb

CatalogNo: YP0334 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA, IHC

MW

- 133kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IF/ICC 1:50-200

ELISA 1:2000-20000

IHC 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human SHIP1 around the phosphorylation site of Tyr1021. AA range:987-1036

Specificity

Phospho-SHIP-1 (Y1021) Polyclonal Antibody detects endogenous levels of SHIP-1 protein only when phosphorylated at Y1021. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PLYGS

| Target Information

Gene name INPP5D

Protein Name Phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase 1

| Organism | Gene ID | UniProt ID |
|----------|-------------------------|--------------------------|
| Human | 3635 ; | Q92835 ; |
| Mouse | 16331 ; | Q9ES52 ; |
| Rat | 54259 ; | P97573 ; |

Cellular Localization


Cytoplasm . Cell membrane ; Peripheral membrane protein . Membrane raft . Cytoplasm , cytoskeleton . Membrane ; Peripheral membrane protein . Translocates to the plasma membrane when activated , translocation is probably due to different mechanisms depending on the stimulus and cell type. Translocates from the cytoplasm to membrane ruffles in a FCGR3/CD16-dependent manner. Colocalizes with FC-gamma-RIIB receptor (FCGR2B) or FCGR3/CD16 at membrane ruffles. Tyrosine phosphorylation may also participate in membrane localization. .

Tissue specificity Specifically expressed in immune and hematopoietic cells. Expressed in bone marrow and blood cells. Levels vary considerably within this compartment. Present in at least 74% of immature CD34+ cells , whereas within the more mature population of CD33+ cells , it is present in only 10% of cells. Present in the majority of T-cells , while it is present in a minority of B-cells (at protein level) .


Function

Catalytic activity:Phosphatidylinositol 3 ,4 ,5-trisphosphate + H (2) O = phosphatidylinositol 3 ,4-bisphosphate + phosphate. ,Domain:The NPXY sequence motif found in many tyrosine-phosphorylated proteins is required for the specific binding of the PID domain. ,Domain:The SH2 domain interacts with tyrosine phosphorylated forms of proteins such as SHC1 or PTPN11/SHP-2. It competes with that of GRB2 for binding to phosphorylated SHC1 to inhibit the Ras pathway. It is also required for tyrosine phosphorylation. ,enzyme regulation:Activated upon translocation to the sites of synthesis of PtdIns (3 ,4 ,5) P3 in the membrane. ,Function:Phosphatidylinositol (PtdIns) phosphatase that specifically hydrolyzes the 5-phosphate of phosphatidylinositol-3 ,4 ,5-trisphosphate (PtdIns (3 ,4 ,5) P3) to produce PtdIns (3 ,4) P2 , thereby negatively regulating the PI3K (phosphoinositide 3-kinase) pathways. Acts as a negative regulator of B-cell antigen receptor signaling. Mediates signaling from the FC-gamma-RIIB receptor (FCGR2B) , playing a central role in terminating signal transduction from activating immune/hematopoietic cell receptor systems. Acts as a negative regulator of myeloid cell proliferation/survival and chemotaxis , mast cell degranulation , immune cells homeostasis , integrin alpha-IIb/beta-3 signaling in platelets and JNK signaling in B-cells. Regulates proliferation of osteoclast precursors , macrophage programming , phagocytosis and activation and is required for endotoxin tolerance. Involved in the control of cell-cell junctions , CD32a signaling in neutrophils and modulation of EGF-induced phospholipase C activity. Key regulator of neutrophil migration , by governing the formation of the leading edge and polarization required for chemotaxis. Modulates FCGR3/CD16-mediated cytotoxicity in NK cells. Mediates the activin/TGF-beta-induced apoptosis through its Smad-dependent expression. May also hydrolyze PtdIns (1 ,3 ,4 ,5) P4 , and could thus affect the levels of the higher inositol polyphosphates like InsP6. ,PTM:Tyrosine phosphorylated by the members of the SRC family after exposure to a diverse array of extracellular stimuli such as cytokines , growth factors , antibodies , chemokines , integrin ligands and hypertonic and oxidative stress. Phosphorylated upon IgG receptor FCGR2B-binding. ,similarity:Belongs to the inositol-1 ,4 ,5-trisphosphate 5-phosphatase family. ,similarity:Contains 1 SH2 domain. ,subcellular location:Translocates to the plasma membrane when activated , translocation is probably due to different mechanisms depending on the stimulus and cell type. Partly translocated via its SH2 domain which mediates interaction with tyrosine phosphorylated receptors such as the FC-gamma-RIIB receptor (FCGR2B) or CD16/FCGR3. Tyrosine phosphorylation may also participate to membrane localization. ,subunit:Interacts with tyrosine phosphorylated forms of SHC1 , DOK1 , DOK3 , PTPN11/SHP-2 , SLAMF1/CD150. Interacts with PTPN11 in response to IL-3. Interacts with receptors EPOR , MS4A2/FCER1B and FCER1G , FCGR2A , FCGR2B and FCGR3. Interacts with GRB2 and PLCG1. Interacts with tyrosine kinases SRC and TEC. Interacts with FCGR2A , leading to regulate gene expression during the phagocytic process. Interacts with c-Met/MET. ,tissue specificity:Specifically expressed in immune and hematopoietic cells. Expressed in bone marrow and blood cells. Levels vary considerably within this compartment. Present in at least 74% of immature CD34+ cells , whereas within the more mature population of CD33+ cells , it is present in only 10% of cells. Present in the majority of T-cells , while it is present in a minority of B-cells (at protein level) . ,


| Validation Data




Western Blot analysis of various cells using Phospho-SHIP-1 (Y1021) Polyclonal Antibody diluted at 1:500




Western Blot analysis of KB cells using Phospho-SHIP-1 (Y1021) Polyclonal Antibody diluted at 1:500



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using SHIP1 (Phospho-Tyr1021) Antibody



Western blot analysis of lysates from HepG2 cells treated with TNF 200NG/ML 30', using SHIP1 (Phospho-Tyr1021) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SHIP-1 (Phospho Tyr1021) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)