

Smad3 (Phospho Ser204) Rabbit pAb

CatalogNo: YP0363 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 48kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Smad3 around the phosphorylation site of Ser204. AA range:170-219

Specificity Phospho-Smad3 (S204) Polyclonal Antibody detects endogenous levels of Smad3 protein only when phosphorylated at S204. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):AGSPN

Target Information

Gene name SMAD3 MADH3

Protein Name Mothers against decapentaplegic homolog 3


Organism	Gene ID	UniProt ID
Human	4088 ;	P84022 ;
Mouse	17127 ;	Q8BUN5 ;
Rat	25631 ;	P84025 ;

Cellular Localization Cytoplasm . Nucleus . Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation , migrates to the nucleus when complexed with SMAD4 (PubMed:15799969 , PubMed:21145499) . Through the action of the phosphatase PPM1A , released from the SMAD2/SMAD4 complex , and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101 , PubMed:19289081) . Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15601644) . MAPK-mediated phosphorylation appears to have no effect on nuclear import (PubMed:19218245) . PDPK1 prevents its nuclear translocation in response to TGF-beta (PubMed:17327236) . Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm of the inner cell mass at the blastocyst stage (By similarity) . .


Tissue specificity Brain ,Colon carcinoma ,Esophagus tumor ,Pancreas ,Placenta ,Spleen ,Umbilical cord blood

Function Disease:Defects in SMAD3 may be a cause of colorectal cancer (CRC) [MIM:114500]. ,Domain:The MH2 domain is sufficient to carry protein nuclear export. ,Function:Transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinase. SMAD3 is a receptor-regulated SMAD (R-SMAD) . ,PTM:Phosphorylated on serine by TGF-beta and activin type 1 receptor kinases. ,similarity:Belongs to the dwarfin/SMAD family. ,similarity:Contains 1 MH1 (MAD homology 1) domain. ,similarity:Contains 1 MH2 (MAD homology 2) domain. ,subcellular location:In the cytoplasm in the absence of ligand. Migration to the nucleus when complexed with Smad4. ,subunit:Interacts with HGS. Interacts with NEDD4L in response to TGF-beta. Interacts with TTRAP (By similarity) . Interacts with SARA (SMAD anchor for receptor activation) ; form trimers with another SMAD3 and the co-SMAD SMAD4. Interacts with JUN/FOS , vitamin D receptor , homeobox protein TGIF and TGIF2 , PEBP2-alpha C subunit , CREB-binding protein (CBP) , p300 , SKI , SNON , ATF2 , SMURF2 , AIP1 , DACH1 and TGFB1I1. Part of a complex consisting of AIP1 , ACVR2A , ACVR1B and SMAD3. Found in a complex with SMAD2 and TRIM33 upon addition of TGF-beta. Interacts with SMAD2 and TRIM33. Found in a complex with SMAD3 , Ran and XPO4. Interacts with XPO4. Interacts with LBXCOR1 and CORL2. ,


Validation Data




Immunofluorescence analysis of Hela cell. 1, mouse Tubulin Antibody YM3035 (green) was diluted at 1:200 (4°C overnight). Smad3 (Phospho Ser204) Rabbit pAb (red) was diluted at 1:200 (4°C overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000 (room temperature, 50min).




Western Blot analysis of various cells using Phospho-Smad3 (S204) Polyclonal Antibody diluted at 1:500



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Smad3 (Phospho-Ser204) Antibody



Western blot analysis of lysates from NIH/3T3 cells treated with Serum 20% 15', using Smad3 (Phospho-Ser204) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Smad3 (Phospho Ser204) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)