

PDCD4 (Phospho Ser67) Rabbit pAb

CatalogNo: YP0387

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat, Monkey

Applications

- WB, IHC, IF, ELISA

MW

- 56kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:10000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human PDCD4 around the phosphorylation site of Ser67. AA range:33-82

Specificity

Phospho-Pdcd-4 (S67) Polyclonal Antibody detects endogenous levels of Pdcd-4 protein only when phosphorylated at S67. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):KNsSR

Target Information

Gene name PDCD4

Protein Name Programmed cell death protein 4


| Organism | Gene ID | UniProt ID |
|----------|-------------------------|--------------------------|
| Human | 27250 ; | Q53EL6 ; |
| Mouse | 18569 ; | Q61823 ; |
| Rat | 64031 ; | Q9JID1 ; |

Cellular Localization Nucleus . Cytoplasm . Shuttles between the nucleus and cytoplasm (By similarity) . Predominantly nuclear under normal growth conditions , and when phosphorylated at Ser-457 (PubMed:16357133) . .


Tissue specificity Up-regulated in proliferative cells. Highly expressed in epithelial cells of the mammary gland. Reduced expression in lung cancer and colon carcinoma.

Function Caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data. ,Disease:Loss of expression correlated with tumor progression of lung and colon carcinoma. ,Domain:Binds EIF4A1 via the MA3 domains. ,Function:Tumor suppressor. Inhibits tumor promoter-induced neoplastic transformation. Down-regulates the expression of MAP4K1 , thus inhibiting events important in driving invasion , namely , MAPK85 activation and consequent JUN-dependent transcription. May play a role in apoptosis. Inhibits the helicase activity of EIF4A and cap-dependent translation. Binds RNA. ,induction:IL2 stimulation inhibits expression , while IL12 increases expression. ,sequence Caution:Contaminating sequence. Potential poly-A sequence. ,similarity:Belongs to the PDCD4 family. ,similarity:Contains 2 MI domains. ,subcellular location:Shuttles between the nucleus and cytoplasm. Predominantly nuclear under normal growth conditions. Exported from the nucleus in the absence of serum. ,subunit:Interacts with EIF4A1 and EIF4A2. ,tissue specificity:Up-regulated in proliferative cells. Highly expressed in epithelial cells of the mammary gland. ,


Validation Data



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from HUVEC cells, K562 cells and COS-7 cells, using PDCD4 (Phospho-Ser67) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
PDCD4 (Phospho Ser67) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)