

TRF1 (Phospho Ser219) Rabbit pAb

CatalogNo: YP0400 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 55kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

ELISA 1:5000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Telomeric Repeat Binding Factor 1 around the phosphorylation site of Ser219. AA range:185-234

Specificity Phospho-TRF1 (S219) Polyclonal Antibody detects endogenous levels of TRF1 protein only when phosphorylated at S219. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):llsQK

| Target Information

Gene name TERF1

Protein Name Telomeric repeat-binding factor 1


Organism	Gene ID	UniProt ID
Human	7013 ;	P54274 ;
Mouse	21749 ;	P70371 ;

Cellular Localization Nucleus. Cytoplasm, cytoskeleton, spindle. Chromosome, telomere. Colocalizes with telomeric DNA in interphase and prophase cells. Telomeric localization decreases in metaphase, anaphase and telophase. Associates with the mitotic spindle.


Tissue specificity Highly expressed and ubiquitous. Isoform Pin2 predominates.

Function Domain:The acidic N-terminal domain binds to the ankyrin repeats of TNKS1 and TNKS2. The C-terminal domain binds microtubules.,Function:Binds the telomeric double-stranded TTAGGG repeat and negatively regulates telomere length. Involved in the regulation of the mitotic spindle. Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of double-stranded TTAGGG repeats added by telomerase and protects chromosome ends; without its protective activity, telomeres are no longer hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways.,induction:Pin2 expression is tightly regulated during the cell cycle; levels are low in G1 and S phase and increase during G2 phase and mitosis.,PTM:ADP-ribosylation by TNKS1 or TNKS2 diminishes its ability to bind to telomeric DNA.,PTM:Phosphorylated preferentially on Ser-219 in an ATM-dependent manner in response to ionizing DNA damage.,similarity:Contains 1 HTH myb-type DNA-binding domain.,subcellular location:Colocalizes with telomeric DNA in interphase and metaphase cells and is located at chromosome ends during metaphase. Associates with the mitotic spindle.,subunit:Homodimer; can contain both isoforms. Found in a complex with POT1; TIN2 and TNKS1. Interacts with ATM, TIN2, TNKS1, TNKS2, PINX1, NEK2 and MAPRE1. Component of the shelterin complex (telosome) composed of TERF1, TERF2, TIN2, TERF2IP ACD and POT1.,tissue specificity:Highly expressed and ubiquitous. Isoform Pin2 predominates.,


| Validation Data



Western Blot analysis of 293 cells using Phospho-TRF1 (S219) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Telomeric Repeat Binding Factor 1 (Phospho-Ser219) Antibody



Western blot analysis of lysates from 293 cells treated with paclitaxel 1uM 24h, using Telomeric Repeat Binding Factor 1 (Phospho-Ser219) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
TRF1 (Phospho Ser219) Rabbit pAb