

Cdc25A (Phospho Thr507) Rabbit pAb

CatalogNo: YP0404

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 59kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

ELISA 1:10000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human Cdc25A (phospho Thr507)

Specificity Phospho-Cdc25A (T507) Polyclonal Antibody detects endogenous levels of Cdc25A protein only when phosphorylated at T507. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): SRtWA

Target Information

Gene name CDC25A

Protein Name M-phase inducer phosphatase 1

Organism	Gene ID	UniProt ID
Human	993 ;	P30304 ;
Mouse		P48964 ;
Rat	171102 ;	P48965 ;

Cellular Localization intracellular ,nucleus ,nucleoplasm ,cytoplasm ,cytosol ,

Tissue specificity Lymph ,

Function Catalytic activity:Protein tyrosine phosphate + H (2) O = protein tyrosine + phosphate. ,Domain:The phosphodegron motif mediates interaction with specific F-box proteins when phosphorylated. Putative phosphorylation sites at Ser-79 and Ser-82 appear to be essential for this interaction. ,enzyme regulation:Stimulated by B-type cyclins. ,Function:Tyrosine protein phosphatase which functions as a dosage-dependent inducer of mitotic progression. Directly dephosphorylates CDC2 and stimulates its kinase activity. Also dephosphorylates CDK2 in complex with cyclin E , in vitro. ,PTM:Phosphorylated by CHEK1 on Ser-76 , Ser-124 , Ser-178 , Ser-279 , Ser-293 and Thr-507 during checkpoint mediated cell cycle arrest. Also phosphorylated by CHEK2 on Ser-124 , Ser-279 , and Ser-293 during checkpoint mediated cell cycle arrest. Phosphorylation on Ser-178 and Thr-507 creates binding sites for YWHAE/14-3-3 epsilon which inhibits CDC25A. Phosphorylation on Ser-76 , Ser-124 , Ser-178 , Ser-279 and Ser-293 may also promote ubiquitin-dependent proteolysis of CDC25A. ,PTM:Ubiquitinated. Association with the F-box proteins BTRC and FBXW11 targets the protein for ubiquitination by CUL1 and proteolysis by the ubiquitin-dependent proteasome pathway. ,similarity:Belongs to the MPI phosphatase family. ,similarity:Contains 1 rhodanese domain. ,subunit:Interacts with CCNB1/cyclin B1. Interacts with YWHAE/14-3-3 epsilon when phosphorylated. Interacts with CUL1 specifically when CUL1 is neddylated and active. Interacts with BTRC/BTRCP1 and FBXW11/BTRCP2. Interactions with CUL1 , BTRC and FBXW11 are enhanced upon DNA damage. ,

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Cdc25A (Phospho Thr507) Rabbit pAb

