

## MAK (Phospho Tyr159) Rabbit pAb

CatalogNo: YP0423 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA

#### MW

- 85kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**ELISA 1:40000**

**Not yet tested in other applications.**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human MAK around the phosphorylation site of Tyr159. AA range:126-175

**Specificity** Phospho-MAK (Y159) Polyclonal Antibody detects endogenous levels of MAK protein only when phosphorylated at Y159. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TDyVS

## Target Information

**Gene name** MAK

**Protein Name** Serine/threonine-protein kinase MAK

Organism	Gene ID	UniProt ID
Human	<a href="#">4117</a> ;	<a href="#">P20794</a> ;
Mouse	<a href="#">17152</a> ;	<a href="#">Q04859</a> ;
Rat	<a href="#">25677</a> ;	<a href="#">P20793</a> ;

**Cellular Localization**

Nucleus. Cytoplasm , cytoskeleton , microtubule organizing center , centrosome. Cytoplasm , cytoskeleton , spindle. Midbody. Cell projection , cilium , photoreceptor outer segment . Photoreceptor inner segment. Localized in both the connecting cilia and the outer segment axonemes (By similarity) . Localized uniformly in nuclei during interphase , to the mitotic spindle and centrosomes during metaphase and anaphase , and also to midbody at anaphase until telophase. .


**Tissue specificity**

Expressed in prostate cancer cell lines at generally higher levels than in normal prostate epithelial cell lines. Isoform 1 is expressed in kidney , testis , lung , trachea , and retina. Isoform 2 is retina-specific where it is expressed in rod and cone photoreceptors.

**Function**

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Could play an important function in spermatogenesis. ,similarity:Belongs to the protein kinase superfamily. ,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily. ,similarity:Contains 1 protein kinase domain. ,tissue specificity:Expressed mainly in testicular cells at and after meiosis. ,

## Validation Data



Western blot analysis of lysates from HepG2 cells treated with PMA 125ng/ml 30', using MAK (Phospho-Tyr159) Antibody. The lane on the right is blocked with the phospho peptide.

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**MAK (Phospho Tyr159) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)