

HP1 γ (Phospho Ser93) Rabbit pAb

CatalogNo: YP0450 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 24kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:10000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human HP1 gamma around the phosphorylation site of Ser93. AA range:59-108

Specificity

Phospho-HP1 γ (S93) Polyclonal Antibody detects endogenous levels of HP1 γ protein only when phosphorylated at S93. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RKSLs

Target Information

Gene name CBX3

Protein Name Chromobox protein homolog 3


Organism	Gene ID	UniProt ID
Human	11335 ; 653972 ;	Q13185 ;
Mouse		P23198 ;

Cellular Localization Nucleus . Associates with euchromatin and is largely excluded from constitutive heterochromatin. May be associated with microtubules and mitotic poles during mitosis (Potential) . .


Tissue specificity Bone marrow ,Brain ,Cajal-Retzius cell ,Epithelium ,Liver ,Placenta ,

Function Function:Seems to be involved in transcriptional silencing in heterochromatin-like complexes. Recognizes and binds histone H3 tails methylated at 'Lys-9' , leading to epigenetic repression. May contribute to the association of the heterochromatin with the inner nuclear membrane through its interaction with lamin B receptor (LBR) . Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins. ,PTM:Phosphorylated by PIM1. Phosphorylated during interphase and possibly hyperphosphorylated during mitosis. ,similarity:Contains 2 chromo domains. ,subcellular location:Associates with euchromatin and is largely excluded from constitutive heterochromatin. May be associated with microtubules and mitotic poles during mitosis. ,subunit:Binds directly to CHAF1A. Interacts with histone H3 methylated at 'Lys-9'. Part of the E2F6.com-1 complex in G0 phase composed of E2F6 , MGA , MAX , TFDP1 , CBX3 , BAT8 , EUHMTASE1 , RING1 , RNF2 , MBLR , L3MBTL2 and YAF2. Interacts with LBR , INCENP , TRIM28/TIF1B , SUV420H1 , SUV420H2 and SP100. Interacts with TIF1A (By similarity) . Interacts with MIS12 and C20orf127. Can interact directly with CBX5 via the chromoshadow domain. ,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HP1 gamma (Phospho-Ser93) Antibody



Western blot analysis of lysates from K562 cells treated with forskolin 40nM 30', using HP1 gamma (Phospho-Ser93) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
HP1 γ (Phospho Ser93) Rabbit pAb