

## Caveolin-2 (Phospho Tyr27) Rabbit pAb

CatalogNo: YP0454 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA

#### MW

- 26kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**ELISA 1:10000**

**Not yet tested in other applications.**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human Caveolin 2 around the phosphorylation site of Tyr27. AA range:12-61

**Specificity** Phospho-Caveolin-2 (Y27) Polyclonal Antibody detects endogenous levels of Caveolin-2 protein only when phosphorylated at Y27. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LEyAD

## Target Information

**Gene name** CAV2

**Protein Name** Caveolin-2

Organism	Gene ID	UniProt ID
Human	<a href="#">858;</a>	<a href="#">P51636;</a>
Mouse	<a href="#">12390;</a>	<a href="#">Q9WVC3;</a>
Rat	<a href="#">363425;</a>	<a href="#">Q2IBC5;</a>

### Cellular Localization


Nucleus. Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Cell membrane; Peripheral membrane protein. Membrane , caveola; Peripheral membrane protein. Potential hairpin-like structure in the membrane. Membrane protein of caveolae. Tyr-19-phosphorylated form is enriched at sites of cell-cell contact and is translocated to the nucleus in complex with MAPK1 in response to insulin (By similarity) . Tyr-27-phosphorylated form is located both in the cytoplasm and plasma membrane. CAV1-mediated Ser-23-phosphorylated form locates to the plasma membrane. Ser-36-phosphorylated form resides in intracellular compartments. .

**Tissue specificity** Expressed in endothelial cells , smooth muscle cells , skeletal myoblasts and fibroblasts.

### Function

Function:May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity. ,Function:May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity. Caveolin-2 may function as an accessory protein in conjunction with caveolin-1. ,online information:Caveolin entry ,similarity:Belongs to the caveolin family. ,subcellular location:Potential hairpin-like structure in the membrane. Membrane protein of caveolae. ,subunit:Homodimer. Caveolin-1 and -2 colocalize and form a stable hetero-oligomeric complex. ,tissue specificity:Expressed in endothelial cells , smooth muscle cells , skeletal myoblasts and fibroblasts. ,

## Validation Data



Western blot analysis of lysates from HepG2 cells treated with EGF 200ng/ml 5' , using Caveolin 2 (Phospho-Tyr27) Antibody. The lane on the right is blocked with the phospho peptide.

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**Caveolin-2 (Phospho Tyr27) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)