

c-Fms (Phospho Tyr699) Rabbit pAb

CatalogNo: YP0504

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 130-170kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

ELISA 1:10000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human c-Fms (phospho Tyr699)

Specificity Phospho-c-Fms (Y699) Polyclonal Antibody detects endogenous levels of c-Fms protein only when phosphorylated at Y699. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):VDyKN

| Target Information

Gene name CSF1R

Protein Name Macrophage colony-stimulating factor 1 receptor

Organism	Gene ID	UniProt ID
Human	1436 ;	P07333 ;
Mouse		P09581 ;

Cellular Localization Cell membrane; Single-pass type I membrane protein.

Tissue specificity Expressed in bone marrow and in differentiated blood mononuclear cells.

Function Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Function:Protein tyrosine-kinase transmembrane receptor for CSF1 and IL34. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 5 Ig-like C2-type (immunoglobulin-like) domains. ,subunit:Interacts with INPPL1/SHIP2 and THOC5. ,tissue specificity:Expressed in bone marrow and in differentiated blood mononuclear cells. ,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
c-Fms (Phospho Tyr699) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)