

## EphA2/5 (Phospho Tyr594) Rabbit pAb

CatalogNo: YP0506

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB, ELISA

#### MW

- 110kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**ELISA 1:10000**

**Not yet tested in other applications.**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized phospho-peptide around the phosphorylation site of human EphA2/5 (phospho Tyr594)

**Specificity** Phospho-EphA2/5 (Y594) Polyclonal Antibody detects endogenous levels of EphA2/5 protein only when phosphorylated at Y594. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):HTyED

## Target Information

**Gene name** EPHA2/EPHA5

**Protein Name** Ephrin type-A receptor 2/5

| Organism | Gene ID   | UniProt ID  |
|----------|---|---|
| Human    | <a href="#">1969</a> ; <a href="#">2044</a> ;   | <a href="#">P29317</a> ; <a href="#">P54756</a> ; |
| Mouse    | <a href="#">13836</a> ; <a href="#">13839</a> ; |   |

### Cellular Localization


Cell membrane ; Single-pass type I membrane protein . Cell projection , ruffle membrane ; Single-pass type I membrane protein . Cell projection , lamellipodium membrane ; Single-pass type I membrane protein . Cell junction , focal adhesion . Present at regions of cell-cell contacts but also at the leading edge of migrating cells (PubMed:19573808 , PubMed:20861311) . Relocates from the plasma membrane to the cytoplasmic and perinuclear regions in cancer cells (PubMed:18794797) . .

**Tissue specificity** Expressed in brain and glioma tissue and glioma cell lines (at protein level) . Expressed most highly in tissues that contain a high proportion of epithelial cells , e.g. skin , intestine , lung , and ovary .

### Function

Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Function:Receptor for members of the ephrin-A family. Binds to ephrin-A1 , -A3 , -A4 and -A5. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 SAM (sterile alpha motif) domain. ,similarity:Contains 2 fibronectin type-III domains. ,subunit:Interacts with SLA (By similarity) . Interacts with INPPL1/SHIP2. ,tissue specificity:Expressed most highly in tissues that contain a high proportion of epithelial cells , e.g. , skin , intestine , lung , and ovary. ,

## Validation Data



Western Blot analysis of JK cells using Phospho-EphA2/5 (Y594) Polyclonal Antibody

## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: http://www.immunoway.com.cn  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**EphA2/5 (Phospho  
Tyr594) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)