

CD45 (Phospho Ser1007) Rabbit pAb

CatalogNo: YP0543 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA

MW

- 150kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IF 1:200-1:1000

ELISA 1:5000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human CD45 around the phosphorylation site of Ser1007. AA range:981-1030

Specificity

Phospho-CD45 (S1007) Polyclonal Antibody detects endogenous levels of CD45 protein only when phosphorylated at S1007. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):DDSDS

Target Information

Gene name PTPRC

Protein Name Receptor-type tyrosine-protein phosphatase C


Organism	Gene ID	UniProt ID
Human	5788 ;	P08575 ;
Mouse		P06800 ;
Rat	24699 ;	P04157 ;

Cellular Localization Cell membrane ; Single-pass type I membrane protein . Membrane raft . Colocalized with DPP4 in membrane rafts. .


Tissue specificity Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3: Detected in thymocytes. Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes. Isoform 7: Detected in thymocytes. Isoform 8: Not detected in thymocytes.

Function Alternative products:At least 8 isoforms are produced ,Catalytic activity:Protein tyrosine phosphate + H (2) O = protein tyrosine + phosphate. ,Disease:Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T (-) B (+) NK (+) SCID) [MIM:608971]. SCID refers to a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity , leukopenia , and low or absent antibody levels. Patients with SCID present in infancy with recurrent , persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. ,Disease:Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain. ,Domain:The first PTPase domain interacts with SKAP1. ,Function:Required for T-cell activation through the antigen receptor. The first PTPase domain has enzymatic activity , while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation , recruits an dephosphorylates SKAP1 and FYN. ,online information:CD45 entry ,online information:PTPRC mutation db ,PTM:Heavily N- and O-glycosylated. ,similarity:Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. ,similarity:Contains 2 fibronectin type-III domains. ,similarity:Contains 2 tyrosine-protein phosphatase domains. ,subunit:Binds GANAB and PRKCSH (By similarity) . Interacts with SKAP1. ,

Validation Data



Immunofluorescence analysis of HeLa cells, using CD45 (Phospho-Ser1007) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with TNF 20ng/ml 15', using CD45 (Phospho-Ser1007) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
CD45 (Phospho Ser1007) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)