

Ksr-1 (Phospho Ser392) Rabbit pAb

CatalogNo: YP0603

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 115kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human KSR around the phosphorylation site of Ser392. AA range:358-407

Specificity

Phospho-Ksr-1 (S392) Polyclonal Antibody detects endogenous levels of Ksr-1 protein only when phosphorylated at S392. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TESVP

Target Information

Gene name KSR1

Protein Name Kinase suppressor of Ras 1

Organism	Gene ID	UniProt ID
Human	8844;	Q8IVT5;
Mouse	16706;	Q61097;

Cellular Localization


Cytoplasm . Membrane ; Peripheral membrane protein . Cell membrane ; Peripheral membrane protein . Cell projection , ruffle membrane . Endoplasmic reticulum membrane . In unstimulated cells , where the phosphorylated form is bound to a 14-3-3 protein , sequestration in the cytoplasm occurs. Following growth factor treatment , the protein is free for membrane translocation , and it moves from the cytoplasm to the cell periphery. .

Tissue specificity Brain ,Epithelium ,Platelet ,Synovial membrane ,


Function

Caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data. ,Function:Location-regulated scaffolding protein connecting MEK to RAF. Promotes MEK and RAF phosphorylation and activity through assembly of an activated signaling complex. By itself , it has no demonstrated kinase activity. ,PTM:Phosphorylated on Ser-309 and , to a higher extent , on Ser-404 by MARK3. Dephosphorylated on Ser-404 by PPP2CA. In resting cells , phosphorylated KSR1 is cytoplasmic and in stimulated cells , dephosphorylated KSR1 is membrane-associated. ,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. ,similarity:Contains 1 phorbol-ester/DAG-type zinc finger. ,similarity:Contains 1 protein kinase domain. ,subcellular location:In unstimulated cells , where the phosphorylated form is bound to a 14-3-3 protein , sequestration in the cytoplasm occurs. Following growth factor treatment , the protein is free for membrane translocation , and it moves from the cytoplasm to the cell periphery. ,subunit:Interacts with HSPCA/HSP90 , YWHAB/14-3-3 , CDC37 , MAP2K/MEK , MARK3 , PPP2R1A and PPP2CA. Also interacts with RAF and MAPK/ERK , in a Ras-dependent manner (By similarity) . The binding of 14-3-3 proteins to phosphorylated KSR prevents the membrane localization. ,


Validation Data




Western Blot analysis of various cells using Phospho-Ksr-1 (S392) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using KSR (Phospho-Ser392) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using KSR (Phospho-Ser392) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HepG2 cells, using KSR (Phospho-Ser392) Antibody. The lane on the right is blocked with the phospho peptide.

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Ksr-1 (Phospho Ser392) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)