

PLC γ 2 (Phospho Tyr753) Rabbit pAb

CatalogNo: YP0608 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 148kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human PLCG2 around the phosphorylation site of Tyr753. AA range:721-770

Specificity

Phospho-PLC γ 2 (Y753) Polyclonal Antibody detects endogenous levels of PLC γ 2 protein only when phosphorylated at Y753. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SLyDV

Target Information

Gene name PLCG2

Protein Name 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-2


Organism	Gene ID	UniProt ID
Human	5336 ;	P16885 ;
Mouse	234779 ;	Q8CIH5 ;
Rat	29337 ;	P24135 ;

Cellular Localization intracellular ,cytosol ,plasma membrane ,extracellular exosome ,


Tissue specificity Lymph ,Lymphoblast ,Spleen ,T-cell ,

Function Catalytic activity:1-phosphatidyl-1D-myo-inositol 4 ,5-bisphosphate + H (2) O = 1D-myo-inositol 1 ,4 ,5-trisphosphate + diacylglycerol. ,cofactor:Calcium. ,Function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1 ,4 ,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling. ,PTM:Phosphorylated on tyrosine residues; upon ligand-induced activation of a variety of growth factor receptors and immune system receptors. Increases phospholipase activity. ,similarity:Contains 1 C2 domain. ,similarity:Contains 1 PH domain. ,similarity:Contains 1 PI-PLC X-box domain. ,similarity:Contains 1 PI-PLC Y-box domain. ,similarity:Contains 1 SH3 domain. ,similarity:Contains 2 SH2 domains. ,


Validation Data




Western Blot analysis of various cells using Phospho-PLC γ 2 (Y753) Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human colon cancer. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using PLCG2 (Phospho-Tyr753) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HepG2 cells treated with Na₃VO₄ 0.3mM 40', using PLCG2 (Phospho-Tyr753) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**PLC γ 2 (Phospho
Tyr753) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)