

## GRK 2 (Phospho Ser29) Rabbit pAb

CatalogNo: YP0653 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 80kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:10000**

**IF 1:50-200**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human GRK2 around the phosphorylation site of Ser29. AA range:14-63

## Specificity

Phospho-GRK 2 (S29) Polyclonal Antibody detects endogenous levels of GRK 2 protein only when phosphorylated at S29. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RASKK

## Target Information

**Gene name** ADRBK1 BARK BARK1 GRK2

**Protein Name** Beta-adrenergic receptor kinase 1


Organism	Gene ID	UniProt ID
Human	<a href="#">156</a> ;	<a href="#">P25098</a> ;
Mouse		<a href="#">Q99MK8</a> ;
Rat	<a href="#">25238</a> ;	<a href="#">P26817</a> ;

**Cellular Localization** Cytoplasm . Cell membrane . Cell junction , synapse , postsynapse . Cell junction , synapse , presynapse .


**Tissue specificity** Expressed in peripheral blood leukocytes.

**Function** Catalytic activity:ATP + [beta-adrenergic receptor] = ADP + [beta-adrenergic receptor] phosphate. ,Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors , probably inducing a desensitization of them. ,online information:Beta adrenergic receptor kinase entry ,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. GPRK subfamily. ,similarity:Contains 1 AGC-kinase C-terminal domain. ,similarity:Contains 1 PH domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 RGS domain. ,subunit:Interacts with GIT1 (By similarity) . Interacts with , and phosphorylates chemokine-stimulated CCR5. ,tissue specificity:Expressed in peripheral blood leukocytes. ,


## Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GRK2 (Phospho-Ser29) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using GRK2 (Phospho-Ser29) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells treated with EGF 200ng/ml 30', using GRK2 (Phospho-Ser29) Antibody. The lane on the right is blocked with the phospho peptide.

## Contact information

Orders: order.cn@immunoway.com  
 Support: support.cn@immunoway.com  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**GRK 2 (Phospho Ser29) Rabbit pAb**

