

www.immunoway.com.cn

NMDAR1 (Phospho Ser896) Rabbit pAb

CatalogNo: YP0696 Orthogonal Validated 💽

Key Features

Host Species Rabbit 	Reactivity • Human,Mouse,Rat	Applications • WB,IHC,IF,ELISA
MW • 105kD (Observed)	Isotype • IgG	

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000 IF 1:50-200

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human NMDAR1
around the phosphorylation site of Ser896. AA range:862-911

Specificity

Phospho-NMDAR1 (S896) Polyclonal Antibody detects endogenous levels of NMDAR1 protein only when phosphorylated at S896. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RRsSK

Target Information

Gene name GRIN1

Protein Name Glutamate [NMDA] receptor subunit zeta-1


Organism	Gene ID	UniProt ID
Human	<u>2902;</u>	<u>Q05586;</u>
Mouse	<u>14810;</u>	<u>P35438;</u>
Rat	<u>24408;</u>	<u>P35439;</u>

CellularCell membrane ; Multi-pass membrane protein . Cell junction, synapse, postsynaptic cell
membrane . Cell junction, synapse, postsynaptic density . Enriched in postsynaptic plasma
membrane and postsynaptic densities. .


Tissue specificity Brain, Cerebellum, Hippocampus,

Function Function:NMDA receptor subtype of glutamate-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Mediated by glycine. This protein plays a key role in synaptic plasticity, synaptogenesis, excitotoxicity, memory acquisition and learning. It mediates neuronal functions in glutamate neurotransmission. Is involved in the cell surface targeting of NMDA receptors.,online information:NMDA receptor entry,PTM:NMDA is probably regulated by C-terminal phosphorylation of an isoform of NR1 by PKC. Dephosphorylated on Ser-897 probably by protein phosphatase 2A (PPP2CB). Its phosphorylated state is influenced by the formation of the NMDAR-PPP2CB complex and the NMDAR channel activity.,similarity:Belongs to the glutamate-gated ion channel (TC 1.A.10) family.,subcellular location:Enriched in post-synaptic plasma membrane and post-synaptic densities.,subunit:Forms heteromeric channel of a zeta subunit (GRIN1), a epsilon subunit (GRIN2A, GRIN2B, GRIN2C or GRIN2D) and a third subunit (GRIN3A or GRIN3B); disulfide-linked. Found in a complex with GRIN2A or GRIN2B, GRIN3A or GRIN3B and PPP2CB. Interacts with DLG4 and MPDZ.,


Validation Data




Western Blot analysis of 293T using Phospho-NMDAζ1 (S896) Polyclonal Antibody. Antibody was diluted at 1:500



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using NMDAR1 (Phospho-Ser896) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using NMDAR1 (Phospho-Ser896) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from K562 cells treated with PMA 125ng/ml 30', using NMDAR1 (Phospho-Ser896) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders:order.cn@immunoway.comSupport:support.cn@immunoway.comTelephone:400-8787-807(China)Website:http://www.immunoway.com.cnAddress:2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: NMDAR1 (Phospho Ser896) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents