

PKC α (Phospho Thr638) Rabbit pAb

CatalogNo: YP0703

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 76kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human PKC alpha around the phosphorylation site of Thr638. AA range:606-655

Specificity

Phospho-PKC α (T638) Polyclonal Antibody detects endogenous levels of PKC α protein only when phosphorylated at T638. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):VLtPP

Target Information

Gene name PRKCA

Protein Name Protein kinase C alpha type

Organism	Gene ID	UniProt ID
Human	5578 ;	P17252 ;
Mouse	18750 ;	P20444 ;
Rat		P05696 ;

Cellular Localization Cytoplasm . Cell membrane ; Peripheral membrane protein . Mitochondrion membrane ; Peripheral membrane protein . Nucleus .

Tissue specificity Blood ,Brain ,Epithelium ,Lung ,Platelet ,

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,cofactor:Binds 3 calcium ions per subunit. The ions are bound to the C2 domain. ,Function:PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters , a class of tumor promoters. ,Function:This is a calcium-activated , phospholipid-dependent , serine- and threonine-specific enzyme. May play a role in cell motility by phosphorylating CSPG4. ,similarity:Belongs to the protein kinase superfamily. ,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily. ,similarity:Contains 1 AGC-kinase C-terminal domain. ,similarity:Contains 1 C2 domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 2 phorbol-ester/DAG-type zinc fingers. ,subunit:Interacts with ADAP1/CENTA1 , CSPG4 and PRKCABP. Binds to SDPR in the presence of phosphatidylserine. ,

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: http://www.immunoway.com.cn
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**PKC α (Phospho
Thr638) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)