

ERK 3 (Phospho Ser189) Rabbit pAb

CatalogNo: YP0738

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 90kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human ERK3 around the phosphorylation site of Ser189. AA range:155-204

Specificity

Phospho-ERK 3 (S189) Polyclonal Antibody detects endogenous levels of ERK 3 protein only when phosphorylated at S189. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):HLSEG

Target Information

Gene name MAPK6

Protein Name Mitogen-activated protein kinase 6

| Organism | Gene ID | UniProt ID |
|----------|-------------------------|--------------------------|
| Human | 5597 ; | Q16659 ; |
| Mouse | 50772 ; | Q61532 ; |
| Rat | 58840 ; | P27704 ; |

Cellular Localization


Cytoplasm . Nucleus . Translocates to the cytoplasm following interaction with MAPKAPK5. .

Tissue specificity Highest expression in the skeletal muscle , followed by the brain. Also found in heart , placenta , lung , liver , pancreas , kidney and skin fibroblasts.


Function

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,cofactor:Magnesium. ,Domain:The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases. ,enzyme regulation:Activated by threonine and tyrosine phosphorylation. ,Function:Phosphorylates microtubule-associated protein 2 (MAP2) . May promote entry in the cell cycle. ,PTM:Dually phosphorylated on Thr-626 and Tyr-628 , which activates the enzyme. ,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. ,similarity:Contains 1 protein kinase domain. ,tissue specificity:Highest expression in the skeletal muscle , followed by the brain. Also found in heart , placenta , lung , liver , pancreas , kidney and skin fibroblasts. ,

Validation Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using ERK3 (Phospho-Ser189) Antibody. The picture on the right is blocked with the phosphopeptide.



Western blot analysis of lysates from mouse brain, using ERK3 (Phospho-Ser189) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
ERK 3 (Phospho Ser189) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)