

hnRNP C1/C2 (Phospho Ser260) Rabbit pAb

CatalogNo: YP0750 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 41kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human hnRNP C1/C2 around the phosphorylation site of Ser260. AA range:231-280

Specificity

Phospho-hnRNP C1/C2 (S260) Polyclonal Antibody detects endogenous levels of hnRNP C1/C2 protein only when phosphorylated at S260. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):DDsAE

| Target Information

Gene name HNRNPC

Protein Name Heterogeneous nuclear ribonucleoproteins C1/C2


Organism	Gene ID	UniProt ID
Human	3183 ;	P07910 ;
Mouse	15381 ;	Q9Z204 ;
Rat		P17132 ;

Cellular Localization Nucleus. Component of ribonucleosomes.


Tissue specificity Blood vessels- blood vessel ,Bone marrow ,Brain ,Cajal

Function Function: Binds pre-mRNA and nucleates the assembly of 40S hnRNP particles. Single HNRNPC tetramers bind 230-240 nucleotides. Trimers of HNRNPC tetramers bind 700 nucleotides. May play a role in the early steps of spliceosome assembly and pre-mRNA splicing. Interacts with poly-U tracts in the 3'-UTR or 5'-UTR of mRNA and modulates the stability and the level of translation of bound mRNA molecules. ,PTM: Phosphorylated on Ser-260 and Ser-299 in resting cells. Phosphorylated on Ser-253 and on 1 serine residue in the poly-Ser stretch at position 238 in response to hydrogen peroxide. ,PTM: Sumoylated. Sumoylation reduces affinity for mRNA. ,similarity: Belongs to the RRM HNRPC family. RALY subfamily. ,similarity: Contains 1 RRM (RNA recognition motif) domain. ,subcellular location: Component of ribonucleosomes. ,subunit: Tetramer composed of 3 copies of isoform C1 and 1 copy of isoform C2. Assembly of 3 tetramers with bound pre-mRNA gives rise to a 19S complex that interacts with HNRNPA2B1 tetramers. Component of the 40S hnRNP particle. Identified in the spliceosome C complex , at least composed of AQR , ASCC3L1 , C19orf29 , CDC40 , CDC5L , CRNKL1 , DDX23 , DDX41 , DDX48 , DDX5 , DGCR14 , DHX35 , DHX38 , DHX8 , EFTUD2 , FRG1 , GPATC1 , HNRNPA1 , HNRNPA2B1 , HNRPA3 , HNRNPC , HNRPF , HNRPH1 , HNRPK , HNRPM , HNRNPR , HNRNPU , KIAA1160 , KIAA1604 , LSM2 , LSM3 , MAGOH , MORG1 , PABPC1 , PLRG1 , PNN , PPIE , PPIL1 , PPIL3 , PPWD1 , PRPF19 , PRPF4B , PRPF6 , PRPF8 , RALY , RBM22 , RBM8A , RBMX , SART1 , SF3A1 , SF3A2 , SF3A3 , SF3B1 , SF3B2 , SF3B3 , SFRS1 , SKIV2L2 , SNRPA1 , SNRPB , SNRPB2 , SNRPD1 , SNRPD2 , SNRPD3 , SNRPE , SNRPF , SNRPG , SNW1 , SRRM1 , SRRM2 , SYF2 , SYNCRIP , TFIP11 , THOC4 , U2AF1 , WDR57 , XAB2 and ZCCHC8. ,


Validation Data




Western Blot analysis of 293 cells using Phospho-hnRNP C1/2 (S260) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using hnRNP C1/2 (Phospho-Ser260) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with H2O2 100uM 15', using hnRNP C1/2 (Phospho-Ser260) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
hnRNP C1/C2
(Phospho Ser260)
Rabbit pAb

