

SRF (Phospho Ser77) Rabbit pAb

CatalogNo: YP0775 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 52kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:40000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human SRF around the phosphorylation site of Ser77. AA range:41-90

Specificity

Phospho-SRF (S77) Polyclonal Antibody detects endogenous levels of SRF protein only when phosphorylated at S77. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LYsGS

Target Information

Gene name SRF

Protein Name Serum response factor


Organism	Gene ID	UniProt ID
Human	6722;	P11831;
Mouse	20807;	Q9JM73;

Cellular Localization Nucleus .


Tissue specificity Brain ,Epithelium ,Lung ,Lymph ,

Function Function:SRF is a transcription factor that binds to the serum response element (SRE) , a short sequence of dyad symmetry located 300 bp to the 5' of the site of transcription initiation of some genes (such as FOS) . Required for cardiac differentiation and maturation. ,PTM:Phosphorylated by PRKDC. ,similarity:Contains 1 MADS-box domain. ,subunit:Binds DNA as a multimer , probably a dimer. Interacts with MLLT7/FOXO4 , NKX3A and SSRP1. Interacts with ARID2 and SRFBP1 (By similarity) . Forms complexes with ARID2 , MYOCD , NKX2-5 and SRFBP1. ,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using SRF (Phospho-Ser77) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using SRF (Phospho-Ser77) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells treated with PMA 125ng/ml 30', using SRF (Phospho-Ser77) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SRF (Phospho Ser77) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)