

FER (Phospho Tyr402) Rabbit pAb

CatalogNo: YP0783 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 85kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human FER around the phosphorylation site of Tyr402. AA range:371-420

Specificity

Phospho-Fer (Y402) Polyclonal Antibody detects endogenous levels of Fer protein only when phosphorylated at Y402. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):VNyEE

Target Information

Gene name FER

Protein Name Tyrosine-protein kinase Fer

Organism	Gene ID	UniProt ID
Human	2241;	P16591;
Mouse	14158;	P70451;

Cellular Localization

Cytoplasm. Cytoplasm , cytoskeleton. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Cell junction. Membrane; Peripheral membrane protein; Cytoplasmic side. Nucleus. Cytoplasm , cell cortex. Associated with the chromatin. Detected on microtubules in polarized and motile vascular endothelial cells. Colocalizes with F-actin at the cell cortex. Colocalizes with PECAM1 and CTNND1 at nascent cell-cell contacts.


Tissue specificity

Isoform 1 is detected in normal colon and in fibroblasts (at protein level) . Isoform 3 is detected in normal testis , in colon carcinoma-derived metastases in lung , liver and ovary , and in colon carcinoma and hepato carcinoma cell lines (at protein level) . Isoform 3 is not detected in normal colon or in normal fibroblasts (at protein level) . Widely expressed.


Function

Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Function:Tyrosine kinase of the non-receptor type. Probably performs an important function , perhaps in regulatory processes such as cell cycle control. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Fes/fps subfamily. ,similarity:Contains 1 FCH domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 SH2 domain. ,subcellular location:Associated with the chromatin. ,tissue specificity:Expressed in a variety of lymphoid cell lines. ,

Validation Data



Immunohistochemistry analysis of paraffin-embedded human brain, using FER (Phospho-Tyr402) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells treated with starved 24h, using FER (Phospho-Tyr402) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
FER (Phospho Tyr402) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)