

## SPAK (Phospho Ser309) Rabbit pAb

CatalogNo: YP0799

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 60kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:5000**

**IF 1:50-200**

### Storage

#### Storage\*

-15°C to -25°C/1 year (Do not lower than -25°C)

#### Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

#### Clonality

Polyclonal

### Immunogen Information

#### Immunogen


The antiserum was produced against synthesized peptide derived from human STK39 around the phosphorylation site of Ser311. AA range: 277-326

**Specificity** Phospho-SPAK (S311) Polyclonal Antibody detects endogenous levels of SPAK protein only when phosphorylated at S311. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): GKsFR


## Target Information

Gene name	STK39		
Protein Name	STE20/SPS1-related proline-alanine-rich protein kinase		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">27347</a> ;	<a href="#">Q9UEW8</a> ;
	Mouse	<a href="#">53416</a> ;	<a href="#">Q9Z1W9</a> ;
	Rat	<a href="#">54348</a> ;	<a href="#">O88506</a> ;
Cellular Localization	Cytoplasm . Nucleus . Nucleus when caspase-cleaved. .		
Tissue specificity	Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.		
Function	Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,Domain:PAPA box (proline-alanine repeats) may target the kinase to a specific subcellular location by facilitating interaction with intracellular proteins such as actin or actin-like proteins.,Function:May act as a mediator of stress-activated signals.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.,similarity:Contains 1 protein kinase domain.,subcellular location:Nucleus when caspase-cleaved.,tissue specificity:Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.,		

## Validation Data



Immunohistochemistry analysis of paraffin-embedded human brain, using STK39 (Phospho-Ser311) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from COLO205 cells, using STK39 (Phospho-Ser311) Antibody. The lane on the right is blocked with the phospho peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
 Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

**SPAK (Phospho Ser309) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)