

Glutamate Receptor 1 (Phospho Ser863) Rabbit pAb

CatalogNo: YP0848

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 102kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:40000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human GluR1 around the phosphorylation site of Ser863. AA range:829-878

Specificity

Phospho-Glutamate Receptor 1 (S863) Polyclonal Antibody detects endogenous levels of Glutamate Receptor 1 protein only when phosphorylated at S863. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RNsGA

| Target Information

Gene name GRIA1

Protein Name Glutamate receptor 1

Organism	Gene ID	UniProt ID
Human	2890 ;	P42261 ;
Mouse	14799 ;	P23818 ;
Rat	50592 ;	P19490 ;

Cellular Localization


Cell membrane ; Multi-pass membrane protein . Endoplasmic reticulum membrane ; Multi-pass membrane protein . Cell junction , synapse , postsynaptic cell membrane ; Multi-pass membrane protein . Cell junction , synapse , postsynaptic density membrane ; Multi-pass membrane protein . Cell projection , dendrite . Cell projection , dendritic spine . Early endosome membrane ; Multi-pass membrane protein . Recycling endosome membrane ; Multi-pass membrane protein . Cell junction , synapse , presynapse . Cell junction , synapse . Interaction with CACNG2 , CNIH2 and CNIH3 promotes cell surface expression. Colocalizes with PDLIM4 in early endosomes. Displays a somatodendritic localization and is excluded from axons in neurons (By similarity) . Localized to cone photoreceptor pedicles (By similarity) . .

Tissue specificity Widely expressed in brain.


Function

Function:Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change , leading to the opening of the cation channel , and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state , characterized by the presence of bound agonist. ,miscellaneous:The postsynaptic actions of Glu are mediated by a variety of receptors that are named according to their selective agonists. This receptor binds AMPA (quisqualate) > glutamate > kainate. ,PTM:Palmitoylated. Depalmitoylated upon glutamate stimulation. Cys-603 palmitoylation leads to Golgi retention and decreased cell surface expression. In contrast , Cys-829 palmitoylation does not affect cell surface expression but regulates stimulation-dependent endocytosis. ,similarity:Belongs to the glutamate-gated ion channel (TC 1.A.10) family. ,subcellular location:Interaction with CACNG2 promotes cell surface expression. ,subunit:Homotetramer or heterotetramer of pore-forming glutamate receptor subunits. Tetramers may be formed by the dimerization of dimers. Interacts with DLG1 via its C-terminus. Interacts with CACNG2 , HIP1 and RASGRF2. ,tissue specificity:Widely expressed in brain. ,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GluR1 (Phospho-Ser863) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using GluR1 (Phospho-Ser863) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from MCF-7 cells, using GluR1 (Phospho-Ser863) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Glutamate Receptor 1 (Phospho Ser863) Rabbit pAb

