

www.immunoway.com.cn

HDAC3 (Phospho Ser424) Rabbit pAb

CatalogNo: YP0921 Orthogonal Validated 💽

Key Features

Host Species Rabbit 	Reactivity Human,Mouse,Rat 	Applications WB,IHC,IF,ELISA
MW • 48kD (Observed)	Isotype • IgG	

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:10000 Not yet tested in other applications.

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human HDAC3
around the phosphorylation site of Ser424. AA range:379-428

Specificity

Phospho-HDAC3 (S424) Polyclonal Antibody detects endogenous levels of HDAC3 protein only when phosphorylated at S424.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):KEsDV

Target Information


Gene name	HDAC3		
Protein Name	Histone deacetylase 3 Organism	Gene ID	UniProt ID
	Human	<u>8841;</u>	<u>015379;</u>
	Mouse		<u>088895;</u>
	Rat	<u>84578;</u>	<u>Q6P6W3;</u>

CellularNucleus . Cytoplasm . Cytoplasm, cytosol . Colocalizes with XBP1 and AKT1 in the cytoplasmLocalization(PubMed:25190803). Predominantly expressed in the nucleus in the presence of CCAR2
(PubMed:21030595). .


Tissue specificity Widely expressed.

Function Catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.,Function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor., PTM: Sumoylated in vitro., similarity: Belongs to the histone deacetylase family. Type 1 subfamily., subunit: Interacts with HDAC7 and HDAC9. Forms a heterologous complex at least with YY1. Interacts with DAXX, HDAC10 and DACH1. Found in a complex with NCOR1 and NCOR2. Component of the N-Cor repressor complex, at least composed of NCOR1, NCOR2, HDAC3, TBL1X, TBL1R, CORO2A and GPS2. Interacts with BCOR, MID2A/IHDM3A, NRIP1, PRDM6 and SRY. Interacts with BTBD14B. Interacts with GLIS2 (By similarity). Interacts with CBFA2T3.,tissue specificity:Widely expressed.,


Validation Data



Immunofluorescence analysis of A549 cells, using HDAC3 (Phospho-Ser424) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human skin, using HDAC3 (Phospho-Ser424) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from NIH/3T3 cells, using HDAC3 (Phospho-Ser424) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

- Orders: order.cn@immunoway.com
- Support: support.cn@immunoway.com
- Telephone: 400-8787-807(China)
- Website: http://www.immunoway.com.cn
- Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: HDAC3 (Phospho Ser424) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents