

## HDAC6 (Phospho Ser22) Rabbit pAb

CatalogNo: YP0922 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 160kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**IF 1:200-1:1000**

**ELISA 1:10000**

Not yet tested in other applications.

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human HDAC6 around the phosphorylation site of Ser22. AA range: 7-56

## Specificity

Phospho-HDAC6 (S22) Polyclonal Antibody detects endogenous levels of HDAC6 protein only when phosphorylated at S22. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PQsPP

## Target Information

**Gene name** HDAC6

**Protein Name** Histone deacetylase 6


Organism	Gene ID	UniProt ID
Human	<a href="#">10013</a> ;	<a href="#">Q9UBN7</a> ;
Mouse	<a href="#">15185</a> ;	<a href="#">Q9Z2V5</a> ;

**Cellular Localization** Cytoplasm . Cytoplasm , cytoskeleton . Nucleus . Perikaryon . Cell projection , dendrite . Cell projection , axon . It is mainly cytoplasmic , where it is associated with microtubules. .


**Tissue specificity** Brain ,Epithelium ,Kidney ,Muscle ,Ovary ,Placenta ,

**Function** Catalytic activity:Hydrolysis of an N (6) -acetyl-lysine residue of a histone to yield a deacetylated histone. ,Function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A , H2B , H3 and H4) . Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation , cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes (By similarity) . Plays a central role in microtubule-dependent cell motility via deacetylation of tubulin. ,PTM:Sumoylated in vitro. ,PTM:Ubiquitinated. Its polyubiquitination however does not lead to its degradation. ,similarity:Belongs to the histone deacetylase family. Type 2 subfamily. ,similarity:Contains 1 UBP-type zinc finger. ,subcellular location:It is mainly cytoplasmic , where it is associated with microtubules. ,subunit:Interacts with CBFA2T3 , HDAC11 and SIRT2. Interacts with F-actin. Interacts with BBIP10. ,


## Validation Data




Immunofluorescence analysis of HeLa cell. 1, mouse Histone H3 Antibody YM3808 (red) was diluted at 1:200 (4°C overnight). HDAC6 (Phospho Ser22) Rabbit pAb (green) was diluted at 1:200 (4°C overnight). 2, Goat Anti Rabbit Alexa Fluor 488 Catalog:RS3211 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 594 Catalog:RS3608 was diluted at 1:1000 (room temperature, 50min).




Western Blot analysis of 3T3 cells using Phospho-HDAC6 (S22) Polyclonal Antibody diluted at 1:500




Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HDAC6 (Phospho-Ser22) Antibody



Immunohistochemistry analysis of paraffin-embedded human testis, using HDAC6 (Phospho-Ser22) Antibody. The picture on the right is blocked with the phosphopeptide.



Western blot analysis of lysates from NIH/3T3 cells treated with Anisomycin 25ug/ml 30', using HDAC6 (Phospho-Ser22) Antibody. The lane on the right is blocked with the phospho peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**HDAC6 (Phospho Ser22) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)