

MLC-2 (Phospho Thr17/Ser18) Rabbit pAb

CatalogNo: YP0957

Key Features

Host Species

- Rabbit

Reactivity

- Human,Mouse,Rat

Applications

- IHC,IF,ELISA

MW

- 20kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:10000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human MLC-2 around the phosphorylation site of Thr17 and Ser18. AA range:5-54

Specificity This antibody detects endogenous levels of MLC-2 only when phosphorylated at Thr17 or Ser18, and dually phosphorylated at two sites. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):AtsNV

Target Information

Gene name MYL12A

Protein Name Myosin regulatory light chain 12A

Organism	Gene ID	UniProt ID
Human	10398 ; 10627 ;	P19105 ; O14950 ; P24844 ;
Mouse	67938 ; 98932 ;	
Rat	50685 ; 296313 ;	P18666 ; Q64122 ;

Cellular Localization stress fiber ,cytosol ,brush border ,myosin complex ,myosin II complex ,Z disc ,apical part of cell ,extracellular exosome ,

Tissue specificity Coronary artery ,Epithelium ,Placenta ,Skeletal muscle ,Skin ,Testis ,

Function Function:Myosin regulatory subunit that plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis , receptor capping , and cell locomotion. ,miscellaneous:This chain binds calcium. ,PTM:Phosphorylation increases the actin-activated myosin ATPase activity and thereby regulates the contractile activity. It is required to generate the driving force in the migration of the cells but not necessary for localization of myosin-2 at the leading edge. ,similarity:Contains 3 EF-hand domains. ,subunit:Myosin is an hexamer of 2 heavy chains and 4 light chains. ,

Validation Data

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
MLC-2 (Phospho Thr17/Ser18) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)