

NFATc1 (Phospho Ser294) Rabbit pAb

CatalogNo: YP0958

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- IHC, IF, ELISA

MW

- 101kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:10000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human NFAT2 around the phosphorylation site of Ser294. AA range:261-310

Specificity Phospho-NFATc1 (S294) Polyclonal Antibody detects endogenous levels of NFATc1 protein only when phosphorylated at S294. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RVsVT

| Target Information

Gene name NFATC1

Protein Name Nuclear factor of activated T-cells cytoplasmic 1

| Organism | Gene ID | UniProt ID |
|----------|------------------------|--------------------------|
| Human | 4772 ; | O95644 ; |
| Mouse | | O88942 ; |


Cellular Localization Cytoplasm . Nucleus . Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription (PubMed:16511445) . Nuclear translocation of NFATC1 is enhanced in the presence of TNFSF11. Nuclear translocation is decreased in the presence of FBN1 which can bind and sequester TNFSF11 (By similarity) . .

Tissue specificity Expressed in thymus , peripheral leukocytes as T-cells and spleen. Isoforms A are preferentially expressed in effector T-cells (thymus and peripheral leukocytes) whereas isoforms B and isoforms C are preferentially expressed in naive T-cells (spleen) . Isoforms B are expressed in naive T-cells after first antigen exposure and isoforms A are expressed in effector T-cells after second antigen exposure. Isoforms IA are widely expressed but not detected in liver nor pancreas , neural expression is strongest in corpus callosum. Isoforms IB are expressed mostly in muscle , cerebellum , placenta and thymus , neural expression in fetal and adult brain , strongest in corpus callosum.


Function

Alternative products:Isoform C-alpha and isoform C-beta are the strongest activator of gene transcription , followed by isoform A-alpha and isoform A-beta , whereas isoform B-alpha and isoform B-beta are the weakest. Isoform B-alpha , isoform B-beta , isoform C-alpha and isoform C-beta , both present in T-cells , can modulate their transcriptional activity ,Domain:Isoforms C have a C-terminal part with an additional trans-activation domain , TAD-B , which acts as a transcriptional activator. Isoforms B have a shorter C-terminal part without complete TAD-B which acts as a transcriptional repressor. ,Domain:Rel Similarity Domain (RSD) allows DNA-binding and cooperative interactions with AP1 factors. ,Domain:The N-terminal transactivation domain (TAD-A) binds to and is activated by Cbp/p300. The dephosphorylated form contains two unmasked nuclear localization signals (NLS) , which allow translocation of the protein to the nucleus. ,Function:Plays a role in the inducible expression of cytokine genes in T-cells , especially in the induction of the IL-2 or IL-4 gene transcription. Also controls gene expression in embryonic cardiac cells. Could regulate not only the activation and proliferation but also the differentiation and programmed death of T-lymphocytes as well as lymphoid and non-lymphoid cells. ,induction:Only isoforms A are inducibly expressed in T lymphocytes upon activation of the T-cell receptor (TCR) complex. Induced after co-addition of phorbol 12-myristate 13-acetate (PMA) and ionomycin. Also induced after co-addition of 12-O-tetradecanoylphorbol-13-acetate (TPA) and ionomycin. Weakly induced with PMA , ionomycin and cyclosporin A. ,PTM:Phosphorylated by NFATC-kinase; dephosphorylated by calcineurin. ,similarity:Contains 1 RHD (Rel-like) domain. ,subcellular location:Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription. ,subunit:Member of the multicomponent NFATC transcription complex that consists of at least two components , a pre-existing cytoplasmic component NFATC2 and an inducible nuclear component NFATC1. Other members such as NFATC4 , NFATC3 or members of the activating protein-1 family , MAF , GATA4 and Cbp/p300 can also bind the complex. NFATC proteins bind to DNA as monomers. ,tissue specificity:Expressed in thymus , peripheral leukocytes as T-cells and spleen. Isoforms A are preferentially expressed in effector T-cells (thymus and peripheral leukocytes) whereas isoforms B and isoforms C are preferentially expressed in naive T-cells (spleen) . Isoforms B are expressed in naive T-cells after first antigen exposure and isoforms A are expressed in effector T-cells after second antigen exposure. ,

Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using NFAT2 (Phospho-Ser294) Antibody



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
NFATc1 (Phospho Ser294) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)