

PAK4 (Phospho Ser474) Rabbit pAb

CatalogNo: YP0973

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 64kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human PAK4/5/6 around the phosphorylation site of Ser474. AA range:441-490

Specificity

Phospho-PAK4 (S474) Polyclonal Antibody detects endogenous levels of PAK4 protein only when phosphorylated at S474. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RKSLV

Target Information

Gene name PAK4

Protein Name Serine/threonine-protein kinase PAK 4

Organism	Gene ID	UniProt ID
Human	10298 ;	O96013 ;
Mouse	70584 ;	Q8BTW9 ;

Cellular Localization


Cytoplasm . Seems to shuttle between cytoplasmic compartments depending on the activating effector. For example , can be found on the cell periphery after activation of growth-factor or integrin-mediated signaling pathways. .

Tissue specificity Highest expression in prostate , testis and colon.

Function


Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Activates the JNK pathway. Plays a role in the reorganization of the actin cytoskeleton and in the formation of filopodia. Phosphorylates and inactivates the protein phosphatase SSH1 , leading to increased inhibitory phosphorylation of the actin binding/depolymerizing factor cofilin. Decreased cofilin activity may lead to stabilization of actin filaments. Phosphorylates ARHGEF2. ,PTM:Autophosphorylated on serine residues when activated by CDC42/p21. ,PTM:Phosphorylated on tyrosine residues upon stimulation of FGFR2. ,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily. ,similarity:Contains 1 CRIB domain. ,similarity:Contains 1 protein kinase domain. ,subunit:Interacts with FGFR2 and GRB2 (By similarity) . Interacts tightly with GTP-bound but not GDP-bound CDC42/p21 and weakly with RAC1. Interacts with its substrate ARHGEF2. ,tissue specificity:Highest expression in prostate , testis and colon. ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

Western Blot analysis of various, using primary antibody at 1:1000 dilution.
Secondary antibody (catalog#:RS23920) was diluted at 1:10000



Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**PAK4 (Phospho
Ser474) Rabbit pAb**