

## MAX (Phospho Ser2) Rabbit pAb

CatalogNo: YP1009

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human,Mouse,Rat

#### Applications

- IHC,IF,ELISA

#### MW

- 18kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**IHC 1:100-1:300**

**ELISA 1:5000**

**IF 1:50-200**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human MAX around the phosphorylation site of Ser2. AA range:1-50

**Specificity** Phospho-Max (S2) Polyclonal Antibody detects endogenous levels of Max protein only when phosphorylated at S2. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):MsDND

---

## | Target Information

**Gene name** MAX BHLHD4

**Protein Name** Protein max

Organism	Gene ID	UniProt ID
Human	<a href="#">4149</a> ;	<a href="#">P61244</a> ;
Mouse		<a href="#">P28574</a> ;
Rat	<a href="#">60661</a> ;	<a href="#">P52164</a> ;


**Cellular Localization** Nucleus. Cell projection , dendrite .

**Tissue specificity** High levels found in the brain , heart and lung while lower levels are seen in the liver , kidney and skeletal muscle.

**Function** Alternative products:Additional isoforms seem to exist ,Caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data. ,Function:Transcription regulator. Forms a sequence-specific DNA-binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC-MAX complex is a transcriptional activator , whereas the MAD-MAX complex is a repressor. May repress transcription via the recruitment of a chromatin remodeling complex containing H3-K9 histone methyltransferase activity. ,PTM:Reversible lysine acetylation might regulate the nuclear-cytoplasmic shuttling of specific Max complexes. ,similarity:Contains 1 basic helix-loop-helix (bHLH) domain. ,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Binds DNA as a heterodimer with MYC or MAD. Part of the E2F6.com-1 complex in G0 phase composed of E2F6 , MGA , MAX , TFDP1 , CBX3 , BAT8 , EUHMTASE1 , RING1 , RNF2 , MBLR , L3MBTL2 and YAF2. Interacts with SPAG9. ,tissue specificity:High levels found in the brain , heart and lung while lower levels are seen in the liver , kidney and skeletal muscle. ,

---

## | Validation Data



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°C overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**MAX (Phospho Ser2)  
Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)