

## LIMK-2 (Phospho Ser283) Rabbit pAb

CatalogNo: YP1028

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat, Monkey

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 72kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:5000**

**IF 1:50-200**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human LIMK2 around the phosphorylation site of Ser283. AA range:249-298

## Specificity

Phospho-LIMK-2 (S283) Polyclonal Antibody detects endogenous levels of LIMK-2 protein only when phosphorylated at S283. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RRSLR

## Target Information

**Gene name** LIMK2

**Protein Name** LIM domain kinase 2

Organism	Gene ID	UniProt ID
Human	<a href="#">3985</a> ;	<a href="#">P53671</a> ;
Mouse	<a href="#">16886</a> ;	<a href="#">O54785</a> ;
Rat	<a href="#">29524</a> ;	<a href="#">P53670</a> ;

### Cellular Localization


Cytoplasm , cytoskeleton , spindle . Cytoplasm , cytoskeleton , microtubule organizing center , centrosome . ; [Isoform LIMK2a]: Cytoplasm . Nucleus . ; [Isoform LIMK2b]: Cytoplasm . Cytoplasm , perinuclear region . Nucleus . Mainly present in the cytoplasm and is scarcely translocated to the nucleus. .

**Tissue specificity** Hepatoma ,Lung ,Ovary ,


### Function

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Displays serine/threonine-specific phosphorylation of myelin basic protein and histone (MBP) in vitro. ,PTM:Phosphorylated on serine and/or threonine residues by ROCK1. ,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. ,similarity:Contains 1 PDZ (DHR) domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 2 LIM zinc-binding domains. ,subcellular location:Isoform LIMK2a is distributed in the cytoplasm and the nucleus. ,subcellular location:Isoform LIMK2b occurs mainly in the cytoplasm and is scarcely translocated to the nucleus. ,subunit:Binds ROCK1 and LKAP. Interacts with PARD3. Interacts with NISCH. ,tissue specificity:Highest expression in the placenta; moderate level in liver , lung , kidney , and pancreas. LIMK2a is found to be more abundant than LIMK2b in liver , colon , stomach , and spleen , while in brain , kidney , and placenta LIMK2b is the dominant form. In adult lung , both LIMK2a and LIMK2b is nearly equally observed. ,


## Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using LIMK2 (Phospho-Ser283) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using LIMK2 (Phospho-Ser283) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of LIMK2 (Phospho-Ser283) Antibody. The lane on the right is blocked with the LIMK2 (Phospho-Ser283) peptide.

## Contact information

Orders: order.cn@immunoway.com  
 Support: support.cn@immunoway.com  
 Telephone: 400-8787-807(China)  
 Website: <http://www.immunoway.com.cn>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**LIMK-2 (Phospho Ser283) Rabbit pAb**