

## Rb (Phospho Ser788) Rabbit pAb

CatalogNo: YP1033

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- IHC, IF, ELISA

#### MW

- 106kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**IHC 1:100-1:300**

**ELISA 1:10000**

**IF 1:50-200**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human Retinoblastoma around the phosphorylation site of Ser788. AA range: 756-805

**Specificity** Phospho-Rb (S788) Polyclonal Antibody detects endogenous levels of Rb protein only when phosphorylated at S788. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): PRsPY

---

## | Target Information

**Gene name** RB1

**Protein Name** Retinoblastoma-associated protein

Organism	Gene ID	UniProt ID
Human	<a href="#">5925</a> ;	<a href="#">P06400</a> ;
Mouse	<a href="#">19645</a> ;	<a href="#">P13405</a> ;
Rat	<a href="#">24708</a> ;	<a href="#">P33568</a> ;


**Cellular Localization** Nucleus . During keratinocyte differentiation , acetylation by KAT2B/PCAF is required for nuclear localization. .

**Tissue specificity** Expressed in the retina. Expressed in foreskin keratinocytes (at protein level) (PubMed:20940255) .

**Function** Disease:Defects in RB1 are a cause of bladder cancer [MIM:109800]. ,Disease:Defects in RB1 are a cause of osteogenic sarcoma [MIM:259500]. ,Disease:Defects in RB1 are the cause of childhood cancer retinoblastoma (RB) [MIM:180200]. RB is a congenital malignant tumor that arises from the nuclear layers of the retina. It occurs in about 1:20'000 live births and represents about 2% of childhood malignancies. It is bilateral in about 30% of cases. Although most RB appear sporadically , about 20% are transmitted as an autosomal dominant trait with incomplete penetrance. The diagnosis is usually made before the age of 2 years when strabismus or a gray to yellow reflex from pupil ("cat eye") is investigated. ,Function:Key regulator of entry into cell division that acts as a tumor suppressor. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated , active form of RB1 interacts with E2F1 and represses its transcription activity , leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and , in particular , that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1 , SUV420H1 and SUV420H2 , leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. In case of viral infections , interactions with SV40 large T antigen , HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity. ,online information:RB1 mutation db ,online information:Retinoblastoma protein entry ,PTM:Phosphorylated in G1 , thereby releasing E2F1 which is then able to activate cell growth. Dephosphorylated at the late M phase. SV40 large T antigen , HPV E7 and adenovirus E1A bind to the underphosphorylated , active form of pRb. ,similarity:Belongs to the retinoblastoma protein (RB) family. ,subunit:Interacts with ATAD5 (By similarity) . The hypophosphorylated form interacts with and sequesters the E2F1 transcription factor. The unphosphorylated form interacts with ARID3B , KDM5A , SUV39H1 , MJD2A/JHDM3A and THOC1. Interacts with the N-terminal domain of TAF1. Interacts with AATF , DNMT1 , LIN9 , LMNA , SUV420H1 , SUV420H2 , PELP1 and TMPO-alpha. May interact with NDC80. Interacts with EID1 and UBR4. Interacts with ARID4A and KDM5B. Interacts with E4F1. Interacts with adenovirus E1A protein , HPV E7 protein and SV40 large T antigen. ,tissue specificity:Expressed in the retina. ,

---

## | Validation Data



Immunohistochemical analysis of paraffin-embedded human Small intestinal stromal tumor. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4°C overnight).3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

**Rb (Phospho Ser788) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)