

STAT2 (Phospho Tyr631) Rabbit pAb

CatalogNo: YP1034

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, ELISA

MW

- 98kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:40000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human STAT2 around the phosphorylation site of Tyr631. AA range:597-646

Specificity Phospho-STAT2 (Y631) Polyclonal Antibody detects endogenous levels of STAT2 protein only when phosphorylated at Y631. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): QPyTK

Target Information

Gene name STAT2

Protein Name Signal transducer and activator of transcription 2


Organism	Gene ID	UniProt ID
Human	6773 ;	P52630 ;
Mouse		Q9WVL2 ;

Cellular Localization Cytoplasm . Nucleus . Translocated into the nucleus upon activation by IFN-alpha/beta. .


Tissue specificity Human small intestine ,Lung ,

Function Function:Signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta) . Following type I IFN binding to cell surface receptors , Jak kinases (TYK2 and JAK1) are activated , leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize , associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor , that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes , which drive the cell in an antiviral state. ,PTM:Tyrosine phosphorylated in response to IFN-alpha. ,similarity:Belongs to the transcription factor STAT family. ,similarity:Contains 1 SH2 domain. ,subcellular location:Translocated into the nucleus upon activation by IFN-alpha/beta. ,subunit:Interacts with ISGF3G/IRF-9 in the cytoplasm. Heterodimer with STAT1 upon IFN-alpha/beta induced phosphorylation. Interacts with CRSP2 and CRSP6. Interacts with Simian virus 5 protein V and rabies virus phosphoprotein. ,

Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using STAT2 (Phospho-Tyr631) Antibody



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
STAT2 (Phospho Tyr631) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)