

JAK2 (Phospho Tyr119) Rabbit pAb

CatalogNo: YP1046

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, ELISA

MW

- 130kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human JAK2 around the phosphorylation site of Tyr119. AA range:81-130

Specificity Phospho-JAK2 (Y119) Polyclonal Antibody detects endogenous levels of JAK2 protein only when phosphorylated at Y119. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RFyFP

Target Information

Gene name JAK2

Protein Name Tyrosine-protein kinase JAK2

Organism	Gene ID	UniProt ID
Human	3717 ;	O60674 ;
Mouse	16452 ;	Q62120 ;
Rat	24514 ;	Q62689 ;

Cellular Localization Endomembrane system ; Peripheral membrane protein . Cytoplasm . Nucleus .

Tissue specificity Ubiquitously expressed throughout most tissues.

Function Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Disease:Chromosomal aberrations involving JAK2 are found in both chronic and acute forms of eosinophilic , lymphoblastic and myeloid leukemia. Translocation t (8;9) (p22;p24) with PCM1 links the protein kinase domain of JAK2 to the major portion of PCM1. Translocation t (9;12) (p24;p13) with ETV6. ,Disease:Defects in JAK2 are a cause of acute myelogenous leukemia (AML) [MIM:601626]. AML is a malignant disease in which hematopoietic precursors are arrested in an early stage of development. ,Disease:Defects in JAK2 are a cause of susceptibility to Budd-Chiari syndrome [MIM:600880]. Budd-Chiari syndrome is a spectrum of disease states , including anatomic abnormalities and hypercoagulable disorders , resulting in hepatic venous outflow occlusion. Clinical manifestations observed in the majority of patients include hepatomegaly , right upper quadrant pain , and abdominal ascites. ,Disease:Defects in JAK2 are associated with familial myelofibrosis [MIM:254450]. Myelofibrosis with myeloid metaplasia is a myeloproliferative disease with annual incidence of 0.5-1.5 cases per 100 ,000 individuals and age at diagnosis around 60 (an increased prevalence is noted in Ashkenazi Jews) . Clinical manifestations depend on the type of blood cell affected and may include anemia , pallor , splenomegaly , hypermetabolic state , petechiae , ecchymosis , bleeding , lymphadenopathy , hepatomegaly , portal hypertension. ,Disease:Defects in JAK2 are associated with polycythemia vera (PV) [MIM:263300]. PV , the most common form of primary polycythemia , is caused by somatic mutation in a single hematopoietic stem cell leading to clonal hematopoiesis. PV is a myeloproliferative disorder characterized predominantly by erythroid hyperplasia , but also by myeloid leukocytosis , thrombocytosis , and splenomegaly. Familial cases of PV are very rare and usually manifest in elderly patients. ,Disease:Defects in JAK2 gene may be a cause of essential thrombocythemia (ET) [MIM:187950]. ET is characterized by elevated platelet levels due to sustained proliferation of megakaryocytes , and frequently lead to thrombotic and haemorrhagic complications. ,Domain:Possesses two phosphotransferase domains. The second one probably contains the catalytic domain (By similarity) , while the presence of slight differences suggest a different role for domain 1. ,Function:Plays a role in leptin signaling and control of body weight (By similarity) . Tyrosine kinase of the non-receptor type , involved in interleukin-3 and probably interleukin-23 signal transduction. ,PTM:Leptin promotes phosphorylation on tyrosine residues , including phosphorylation on Tyr-813. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. JAK subfamily. ,similarity:Contains 1 FERM domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 SH2 domain. ,subcellular location:Wholly intracellular , possibly membrane associated. ,subunit:Interacts with SIRPA and SH2B1 (By similarity) . Interacts with IL23R , SKB1 and STAM2. ,tissue specificity:Expressed in blood , bone marrow and lymph node. ,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
JAK2 (Phospho Tyr119) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)