

COT (Phospho Ser400) Rabbit pAb

CatalogNo: YP1049

Key Features

Host Species

- Rabbit

Reactivity

- Human,Mouse,Rat

Applications

- IHC,IF,ELISA

MW

- 53kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human MAP3K8 around the phosphorylation site of Ser400. AA range:366-415

Specificity Phospho-Cot (S400) Polyclonal Antibody detects endogenous levels of Cot protein only when phosphorylated at S400. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):CQsLD

Target Information

Gene name MAP3K8

Protein Name Mitogen-activated protein kinase kinase kinase 8


Organism	Gene ID	UniProt ID
Human	1326 ;	P41279 ;
Mouse	26410 ;	Q07174 ;
Rat	116596 ;	Q63562 ;

Cellular Localization Cytoplasm .

Tissue specificity Expressed in several normal tissues and human tumor-derived cell lines.

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,cofactor:Magnesium. ,developmental stage:Isoform 1 is activated specifically during the S and G2/M phases of the cell cycle. ,Function:Required for TLR4 activation of the MEK/ERK pathway. Able to activate NF-kappa-B 1 by stimulating proteasome-mediated proteolysis of NF-kappa-B 1/p105. Plays a role in the cell cycle. The longer form has some transforming activity , although it is much weaker than the activated cot oncoprotein. ,PTM:Autophosphorylated. Isoform 1 undergoes phosphorylation mainly on Ser residues , and isoform 2 on both Ser and Thr residues. ,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily. ,similarity:Contains 1 protein kinase domain. ,subunit:Forms a ternary complex with NFKB1 and TNIP2. ,tissue specificity:Expressed in several normal tissues and human tumor-derived cell lines. ,

Validation Data



Immunohistochemistry analysis of paraffin-embedded human brain, using MAP3K8 (Phospho-Ser400) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
COT (Phospho Ser400) Rabbit pAb

