

## iNOS (Phospho Tyr151) Rabbit pAb

CatalogNo: YP1067

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- IHC, IF, ELISA

#### MW

- 131kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**IHC 1:100-1:300**

**ELISA 1:10000**

**IF 1:50-200**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human iNOS around the phosphorylation site of Tyr151. AA range:117-166

**Specificity** Phospho-NOS2 (Y151) Polyclonal Antibody detects endogenous levels of NOS2 protein only when phosphorylated at Y151. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):QYyGS

## Target Information

**Gene name** NOS2, INOS

**Protein Name** Nitric oxide synthase inducible


Organism	Gene ID	UniProt ID
Human	<a href="#">4843</a> ;	<a href="#">P35228</a> ;
Mouse	<a href="#">18126</a> ;	<a href="#">P29477</a> ;
Rat	<a href="#">24599</a> ;	<a href="#">Q06518</a> ;

**Cellular Localization** Cytoplasm , cytosol . Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4 , exhibits a more diffuse cytosolic localization. .


**Tissue specificity** Expressed in the liver , retina , bone cells and airway epithelial cells of the lung. Not expressed in the platelets. Expressed in chondrocytes (PubMed:7504305) .

**Function** Catalytic activity:L-arginine + n NADPH + n H (+) + m O (2) = citrulline + nitric oxide + n NADP (+) . ,cofactor:Binds 1 FAD. ,cofactor:Binds 1 FMN. ,cofactor:Heme group. ,cofactor:Tetrahydrobiopterin (BH4) . May stabilize the dimeric form of the enzyme. ,enzyme regulation:Regulated by calcium/calmodulin. Aspirin inhibits expression and function of this enzyme and effects may be exerted at the level of translational/post-translational modification and directly on the catalytic activity. ,Function:Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages , NO mediates tumoricidal and bactericidal actions. ,induction:By endotoxins and cytokines. ,online information:Nitric oxide synthase entry ,similarity:Belongs to the NOS family. ,similarity:Contains 1 FAD-binding FR-type domain. ,similarity:Contains 1 flavodoxin-like domain. ,subunit:Homodimer. Binds SLC9A3R1. ,tissue specificity:Expressed in the liver , retina , bone cells and airway epithelial cells of the lung. Not expressed in the platelets. ,

## Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using iNOS (Phospho-Tyr151) Antibody



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using iNOS (Phospho-Tyr151) Antibody. The picture on the right is blocked with the phospho peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**iNOS (Phospho Tyr151) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)