

HP1 α (Phospho Ser92) Rabbit pAb

CatalogNo: YP1073

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, ELISA

MW

- 22kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:40000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human HP1 alpha around the phosphorylation site of Ser92. AA range:58-107

Specificity Phospho-HP1 α (S92) Polyclonal Antibody detects endogenous levels of HP1 α protein only when phosphorylated at S92. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RKsNF

Target Information

Gene name CBX5 HP1A

Protein Name Chromobox protein homolog 5 (Antigen p25) (Heterochromatin protein 1 homolog alpha) (HP1 alpha)


Organism	Gene ID	UniProt ID
Human	23468 ;	P45973 ;
Mouse	12419 ;	Q61686 ;

Cellular Localization Nucleus . Chromosome . Chromosome , centromere . Colocalizes with HNRNPU in the nucleus (PubMed:19617346) . Component of centromeric and pericentromeric heterochromatin. Associates with chromosomes during mitosis. Associates specifically with chromatin during metaphase and anaphase. .


Tissue specificity Epithelium ,Fetal brain cortex ,Placenta ,

Function Function:Component of heterochromatin. Recognizes and binds histone H3 tails methylated at 'Lys-9' , leading to epigenetic repression. Can interact with lamin B receptor (LBR) . This interaction can contribute to the association of the heterochromatin with the inner nuclear membrane. Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins. ,PTM:Phosphorylation of HP1 and LBR may be responsible for some of the alterations in chromatin organization and nuclear structure which occur at various times during the cell cycle (By similarity) . Phosphorylated during interphase and possibly hyper-phosphorylated during mitosis. ,similarity:Contains 2 chromo domains. ,subcellular location:Component of centromeric and pericentromeric heterochromatin. Associates with chromosomes during mitosis. Associates specifically with chromatin during metaphase and anaphase. ,subunit:Interacts with SUV420H1 and SUV420H2 (By similarity) . Interacts directly with ATRX , CHAF1A , LBR , NIPBL , SP100 , STAM2 and TRIM28 via the chromoshadow domain. Can interact directly with CBX3 via the chromoshadow domain. Interacts with histone H3 methylated at 'Lys-9'. Interacts with MIS12 and C20orf127. Interacts with HP1BP3. ,

Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HP1 alpha (Phospho-Ser92) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using HP1 alpha (Phospho-Ser92) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

HP1 α (Phospho Ser92) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)