

ER β (Phospho Ser87) Rabbit pAb

CatalogNo: YP1076

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, ELISA

MW

- 59kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Estrogen Receptor-beta around the phosphorylation site of Ser87. AA range:53-102

Specificity Phospho-ER β (S87) Polyclonal Antibody detects endogenous levels of ER β protein only when phosphorylated at S87. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):HLsPL

| Target Information

Gene name ESR2 ESTRB NR3A2

Protein Name Estrogen receptor beta


Organism	Gene ID	UniProt ID
Human	2100 ;	Q92731 ;
Mouse	13983 ;	O08537 ;
Rat	25149 ;	Q62986 ;

Cellular Localization Nucleus .

Tissue specificity [Isoform 1]: Expressed in testis and ovary , and at a lower level in heart , brain , placenta , liver , skeletal muscle , spleen , thymus , prostate , colon , bone marrow , mammary gland and uterus. Also found in uterine bone , breast , and ovarian tumor cell lines , but not in colon and liver tumors. ; [Isoform 2]: Expressed in spleen , thymus , testis and ovary and at a lower level in skeletal muscle , prostate , colon , small intestine , leukocytes , bone marrow , mammary gland and uterus. ; [Isoform 4]: Expressed in the testis. ; [Isoform 5]: Expressed in testis , and at a lower level in spleen , thymus , ovary , mammary gland and uterus. ; [Isoform 6]: Expressed in testis , placenta , skeletal muscle , spleen and leukocytes , and at a lower level in heart , lung , liver , kidney , pancreas , thymus , prostate , colon , small intestine , bone marrow , mammary gland and uterus. Not expressed in brain.

Function Domain:Composed of three domains: a modulating N-terminal domain , a DNA-binding domain and a C-terminal steroid-binding domain. ,Function:Nuclear hormone receptor. Binds estrogens with an affinity similar to that of ESR1 , and activates expression of reporter genes containing estrogen response elements (ERE) in an estrogen-dependent manner. Isoform beta-cx lacks ligand binding ability and has no or only very low ere binding activity resulting in the loss of ligand-dependent transactivation ability. DNA-binding by ESR1 and ESR2 is rapidly lost at 37 degrees Celsius in the absence of ligand while in the presence of 17 beta-estradiol and 4-hydroxy-tamoxifen loss in DNA-binding at elevated temperature is more gradual. ,online information:Estrogen receptor entry ,similarity:Belongs to the nuclear hormone receptor family. ,similarity:Belongs to the nuclear hormone receptor family. NR3 subfamily. ,similarity:Contains 1 nuclear receptor DNA-binding domain. ,subunit:Binds DNA as a homodimer. Can form a heterodimer with ESR1. Interacts with NCOA3 , NCOA5 and NCOA6 coactivators , leading to a strong increase of transcription of target genes. Interacts with PELP1 and UBE1C. Isoform beta-2/cx preferentially forms a heterodimer with ESR1 rather than ESR2 and inhibits DNA-binding by ESR1. Interacts with AKAP13. Interacts with DNTTIP2. Interacts with isoform 4 of TXNRD1. ,tissue specificity:Isoform beta-1 is expressed in testis and ovary , and at a lower level in heart , brain , placenta , liver , skeletal muscle , spleen , thymus , prostate , colon , bone marrow , mammary gland and uterus. Also found in uterine bone , breast , and ovarian tumor cell lines , but not in colon and liver tumors. Isoform beta-2 is expressed in spleen , thymus , testis and ovary and at a lower level in skeletal muscle , prostate , colon , small intestine , leukocytes , bone marrow , mammary gland and uterus. Isoform beta-3 is found in testis. Isoform beta-4 is expressed in testis , and at a lower level in spleen , thymus , ovary , mammary gland and uterus. Isoform beta-5 is expressed in testis , placenta , skeletal muscle , spleen and leukocytes , and at a lower level in heart , lung , liver , kidney , pancreas , thymus , prostate , colon , small intestine , bone marrow , mammary gland and uterus. Not expressed in brain. ,

| Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
ERβ (Phospho Ser87) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)