

Sgo1 (Phospho Ser14) Rabbit pAb

CatalogNo: YP1107

Key Features

Host Species

- Rabbit

Reactivity

- Human,Rat,Mouse,

Applications

- IHC,IF,ELISA

MW

- 64kD (Calculated)

Isotype

- IgG

Recommended Dilution Ratios

IHC 1:100-1:300

ELISA 1:5000

IF 1:50-200

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

The antiserum was produced against synthesized peptide derived from human SGO1 around the phosphorylation site of Ser14. AA range:1-50

Specificity

Phospho-Sgo1 (S14) Polyclonal Antibody detects endogenous levels of Sgo1 protein only when phosphorylated at S14.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):QDsLE

| Target Information

Gene name SGOL1

Protein Name Shugoshin-like 1


Organism	Gene ID	UniProt ID
Human	151648 ;	Q5FBB7 ;
Mouse		Q9CXH7 ;

Cellular Localization Nucleus . Chromosome, centromere . Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, spindle pole . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Localizes to the inner centromere throughout prophase until metaphase and disappears at anaphase (PubMed:16541025). Centromeric localization requires the presence of BUB1 and the interaction with PPP2R1A (PubMed:16580887)(PubMed:16541025)(PubMed:15604152). Colocalizes with NEK2 at the kinetochore (PubMed:17621308). Colocalizes with and SS18L1 at the kinetochore (PubMed:16582621). Phosphorylation by AUKRB and the presence of BUB1 are required for localization to the kinetochore (PubMed:17617734). Isoform 1 primarily localizes to kinetochores during G2 phase and mitotic prophase, metaphase, and anaphase and does not appear to be associated with kinetochores during late mitosis (PubMed:16582621). Isoform 3 is found at the centrosome in interphase and at spindle poles in mitosis and its spindle pole localization is PLK1 dependent (PubMed:16582621). Isoform 3 does not localize to kinetochores during any stages of the cell cycle (PubMed:16582621). .

Tissue specificity Widely expressed. Highly expressed in testis. Expressed in lung, small intestine, breast, liver and placenta. Strongly overexpressed in 90% of breast cancers tested.

Function developmental stage:Appears in prophase cells and remains present until metaphase. Strongly decreases at the onset of anaphase and completely disappears at telophase. Not present in interphase cells (at protein level).,Domain:The D-box (destruction box) mediates the interaction with APC proteins, and may act as a recognition signal for degradation via the ubiquitin-proteasome pathway.,Function:Plays a central role in chromosome cohesion during mitosis by preventing premature dissociation of cohesin complex from centromeres after prophase, when most of cohesin complex dissociates from chromosomes arms. May act by preventing phosphorylation of the STAG2 subunit of cohesin complex at the centromere, ensuring cohesin persistence at centromere until cohesin cleavage by ESPL1/separase at anaphase.,miscellaneous:Strongly overexpressed in 90% of breast cancers tested.,PTM:Ubiquitinated by the anaphase promoting complex (APC) at the onset of anaphase, conducting to its degradation.,similarity:Belongs to the shugoshin family.,subcellular location:Localizes to the centromere throughout prophase until metaphase and disappears at anaphase. BUB1 is required for centromeric localization. During prometaphase, it localizes to a single focus, while at metaphase, it localizes to 2 spots corresponding to the 2 centromeres.,subunit:Interacts with PPP2CA (or PPP2CB), PPP2R1B, PPP2R5A, PPP2R5B, PPP2R5C, PPP2R5D, PPP2R5E, SET, LRRC59, RBM10 (or RBM5), RPL10A, RPL28, RPL7, RPL7A and RPLP1. Interaction with protein phosphatase 2A occurs most probably through direct binding to the regulatory B56 subunits: PPP2R1B, PPP2R5A, PPP2R5B, PPP2R5C, PPP2R5D, PPP2R5E.,tissue specificity:Widely expressed. Highly expressed in testis. Expressed in lung, small intestine, breast, liver and placenta.,

| Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Sgo1 (Phospho Ser14) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)