

TBC1D4 (Phospho Thr642) Rabbit pAb

CatalogNo: YP1128

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 150kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:5000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human AS160 around the phosphorylation site of Thr642. AA range:611-660

Specificity

Phospho-TBC1D4 (T642) Polyclonal Antibody detects endogenous levels of TBC1D4 protein only when phosphorylated at T642. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):AHTFS

Target Information

Gene name TBC1D4 AS160 KIAA0603

Protein Name TBC1 domain family member 4


Organism	Gene ID	UniProt ID
Human	9882 ;	O60343 ;
Mouse	210789 ;	Q8BYJ6 ;

Cellular Localization Cytoplasm . Isoform 2 shows a cytoplasmic perinuclear localization in a myoblastic cell line in resting and insulin-stimulated cells.


Tissue specificity Widely expressed. Isoform 2 is the highest overexpressed in most tissues. Isoform 1 is highly expressed in skeletal muscle and heart , but was not detectable in the liver nor in adipose tissue. Isoform 2 is strongly expressed in adrenal and thyroid gland , and also in lung , kidney , colon , brain and adipose tissue. Isoform 2 is moderately expressed in skeletal muscle. Expressed in pancreatic Langerhans islets , including beta cells (at protein level) . Expression is decreased by twofold in pancreatic islets in type 2 diabetes patients compared to control subjects. Up-regulated in T-cells from patients with atopic dermatitis.

Function Disease:May be involved in atopic dermatitis (AD) . ,Function:May act as a GTPase-activating protein for RAB2A , RAB8A , RAB10 and RAB14. Isoform 2 promotes insulin-induced glucose transporter SLC2A4/GLUT4 translocation at the plasma membrane , thus increasing glucose uptake. ,PTM:Insulin-stimulated phosphorylation is required for SLC2A4/GLUT4 translocation. ,PTM:Phosphorylated by AKT1; insulin-induced. ,PTM:Physiological hyperinsulinemia increases phosphorylation in skeletal muscle. Insulin-stimulated phosphorylation is reduced by 39% in type 2 diabetic patients. ,similarity:Contains 1 Rab-GAP TBC domain. ,similarity:Contains 2 PID domains. ,subcellular location:Isoform 2 shows a cytoplasmic perinuclear localization in a myoblastic cell line in resting and insulin-stimulated cells. ,tissue specificity:Widely expressed , but differential expression for isoforms 1 and 2 , with highest overall expression of isoform 2 in most tissues. Isoform 1 is highly expressed in skeletal muscle and heart , but was not detectable in the liver nor in adipose tissue. Isoform 2 strongly expressed in adrenal and thyroid gland , and also in lung , kidney , colon , brain and adipose tissue. Moderate isoform 2 expression in skeletal muscle. Expressed in pancreatic Langerhans islets , including beta cells (at protein level) . Expression is decreased by twofold in pancreatic islets in type 2 diabetes patients compared to control subjects. ,


Validation Data



Western blot analysis of L929 using p-TBC1D4 (T642) antibody. Antibody was diluted at 1:2000



Immunofluorescence analysis of HeLa cells, using AS160 (Phospho-Thr642) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using AS160 (Phospho-Thr642) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
TBC1D4 (Phospho Thr642) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)