

## FAK (Phospho Ser910) Rabbit pAb

CatalogNo: YP1146

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- IHC, IF, ELISA

#### MW

- 119kD (Calculated)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**IHC 1:100-1:300**

**IF 1:200-1:1000**

**ELISA 1:5000**

**Not yet tested in other applications.**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human FAK around the phosphorylation site of Ser910. AA range:876-925

**Specificity**

Phospho-FAK (S910) Polyclonal Antibody detects endogenous levels of FAK protein only when phosphorylated at S910. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):EISPP

**| Target Information**

**Gene name** PTK2

**Protein Name** Focal adhesion kinase 1

Organism	Gene ID	UniProt ID
Human	<a href="#">5747;</a>	<a href="#">Q05397;</a>
Mouse	<a href="#">14083;</a>	<a href="#">P34152;</a>
Rat	<a href="#">25614;</a>	<a href="#">O35346;</a>

**Cellular Localization**

Cell junction , focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm , cell cortex. Cytoplasm , cytoskeleton. Cytoplasm , cytoskeleton , microtubule organizing center , centrosome . Nucleus. Cytoplasm , cytoskeleton , cilium basal body . Constituent of focal adhesions. Detected at microtubules.


**Tissue specificity**

Detected in B and T-lymphocytes. Isoform 1 and isoform 6 are detected in lung fibroblasts (at protein level) . Ubiquitous. Expressed in epithelial cells (at protein level) (PubMed:31630787) .


**Function**

Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Domain:The carboxy-terminal region is the site of focal adhesion targeting (FAT) sequence which mediates the localization of FAK1 to focal adhesions. ,Domain:The first Pro-rich domain interacts with the SH3 domain of CRK-associated substrate (BCAR1) and CASL. ,Function:Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility , proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking , or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid , or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity. ,PTM:Phosphorylated on 6 tyrosine residues upon activation. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. FAK subfamily. ,similarity:Contains 1 FERM domain. ,similarity:Contains 1 protein kinase domain. ,subcellular location:Constituent of focal adhesions. ,subunit:Interacts with CAS family members and with GIT1 , SORBS1 and BCAR3. Interacts with RGFNEF and SHB (By similarity) . Interacts with TGFB1I1. ,tissue specificity:Expressed in all organs tested , in lymphoid cell lines , but most abundantly in brain. ,

## Validation Data



Immunofluorescence analysis of HepG2 cells, using FAK (Phospho-Ser910) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using FAK (Phospho-Ser910) Antibody. The picture on the right is blocked with the phospho peptide.

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**FAK (Phospho Ser910) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)