

Synapsin I (Phospho Ser605) Rabbit pAb

CatalogNo: YP1149

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, ELISA

MW

- 74kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Synapsin1 around the phosphorylation site of Ser605. AA range:576-625

Specificity

Phospho-Synapsin I (S605) Polyclonal Antibody detects endogenous levels of Synapsin I protein only when phosphorylated at S605. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): QAsQA

Target Information

Gene name SYN1

Protein Name Synapsin-1


Organism	Gene ID	UniProt ID
Human	6853 ;	P17600 ;
Mouse	20964 ;	O88935 ;
Rat	24949 ;	P09951 ;

Cellular Localization Cell junction , synapse. Golgi apparatus .


Tissue specificity Brain ,Brain cortex ,

Function Disease: Defects in SYN1 are a cause of epilepsy X-linked with variable learning disabilities and behavior disorders [MIM:300491]. XELBD is characterized by variable combinations of epilepsy , learning difficulties , macrocephaly , and aggressive behavior. ,Function: Neuronal phosphoprotein that coats synaptic vesicles , binds to the cytoskeleton , and is believed to function in the regulation of neurotransmitter release. The complex formed with NOS1 and CAPON proteins is necessary for specific nitric-oxid functions at a presynaptic level. ,PTM: Substrate of at least four different protein kinases. It is probable that phosphorylation plays a role in the regulation of synapsin-1 in the nerve terminal. Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity: Belongs to the synapsin family. ,subunit: Homodimer. Interacts with CAPON. Forms a ternary complex with NOS1. Isoform Ib interacts with PRNP. ,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Synapsin1 (Phospho-Ser605) Antibody



Immunofluorescence analysis of COS7 cells, using Synapsin1 (Phospho-Ser605) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using Synapsin1 (Phospho-Ser605) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Synapsin I (Phospho Ser605) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)