

DNA-PKCS (Phospho Thr2638) Rabbit pAb

CatalogNo: YP1174 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- IF, ELISA

MW

- 469kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IF 1:200-1:1000

ELISA 1:10000

Not yet tested in other applications

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human DNA-PK around the phosphorylation site of Thr2638. AA range:2604-2653

Specificity Phospho-DNA-PKCS (T2638) Polyclonal Antibody detects endogenous levels of DNA-PKCS protein only when phosphorylated at T2638. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RATQQ

Target Information

Gene name PRKDC

Protein Name DNA-dependent protein kinase catalytic subunit


Organism	Gene ID	UniProt ID
Human	5591 ;	P78527 ;
Mouse	19090 ;	P97313 ;

Cellular Localization Nucleus . Nucleus , nucleolus .


Tissue specificity Brain ,Cervix carcinoma ,Epithelium ,Fetal lung ,Placen

Function Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,enzyme regulation:Inhibited by wortmannin. Activity of the enzyme seems to be attenuated by autophosphorylation. ,Function:Serine/threonine-protein kinase that acts as a molecular sensor for DNA damage. Involved in DNA nonhomologous end joining (NHEJ) required for double-strand break (DSB) repair and V (D) J recombination. Must be bound to DNA to express its catalytic properties. Promotes processing of hairpin DNA structures in V (D) J recombination by activation of the hairpin endonuclease artemis (DCLRE1C) . The assembly of the DNA-PK complex at DNA ends is also required for the NHEJ ligation step. Required to protect and align broken ends of DNA. May also act as a scaffold protein to aid the localization of DNA repair proteins to the site of damage. Found at the ends of chromosomes , suggesting a further role in the maintenance of telomeric stability and the prevention of chromosomal end fusion. Also involved in modulation of transcription. Recognizes the substrate consensus sequence [ST]-Q. Phosphorylates 'Ser-139' of histone variant H2AX/H2AFX , thereby regulating DNA damage response mechanism. Phosphorylates DCLRE1C , c-Abl/ABL1 , histone H1 , HSPCA , c-jun/JUN , p53/TP53 , PARP1 , POU2F1 , DHX9 , SRF , XRCC1 , XRCC1 , XRCC4 , XRCC5 , XRCC6 , WRN , c-myc/MYC and RFA2. Can phosphorylate C1D not only in the presence of linear DNA but also in the presence of supercoiled DNA. Ability to phosphorylate TP53/p53 in the presence of supercoiled DNA is dependent on C1D. ,PTM:Phosphorylated upon DNA damage , probably by ATM or ATR. Autophosphorylated on Thr-2609 , Thr-2638 and Thr-2647. Thr-2609 is a DNA damage-inducible phosphorylation site (inducible with ionizing radiation , IR) . Autophosphorylation induces a conformational change that leads to remodeling of the DNA-PK complex , requisite for efficient end processing and DNA repair. ,similarity:Belongs to the PI3/PI4-kinase family. ,similarity:Contains 1 FAT domain. ,similarity:Contains 1 FATC domain. ,similarity:Contains 1 PI3K/PI4K domain. ,similarity:Contains 2 HEAT repeats. ,similarity:Contains 3 TPR repeats. ,subunit:DNA-PK is a heterotrimer of PRKDC and the Ku p70-p86 (XRCC6-XRCC5) dimer. Formation of this complex may be promoted by interaction with ILF3. Associates with the DNA-bound Ku heterodimer , but it can also bind to and be activated by free DNA. Interacts with DNA-PKcs-interacting protein (KIP) with the region upstream the kinase domain. PRKDC alone also interacts with and phosphorylates DCLRE1C , thereby activating the latent endonuclease activity of this protein. Interacts with C1D. ,

Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using DNA-PK (Phospho-Thr2638) Antibody



Immunofluorescence analysis of HUVEC cells treated with serum 20% 30', using DNA-PK (Phospho-Thr2638) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
DNA-PKCS (Phospho Thr2638) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)