

Ob-R (Phospho Tyr1141) Rabbit pAb

CatalogNo: YP1200

Key Features

Host Species Rabbit 	Reactivity Human,Mouse 	ApplicationsWB,ELISA
MW • 132kD (Observed)	Isotype • IgG	

Recommended Dilution Ratios

WB 1:500-1:2000 ELISA 1:5000 Not yet tested in other applications.

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenSynthesized phospho-peptide around the phosphorylation site of human Ob-R (phospho
Tyr1141)SpecificityPhospho-Ob-R (Y1141) Polyclonal Antibody detects endogenous levels of Ob-R protein
only when phosphorylated at Y1141.The name of modified sites may be influenced by
many factors, such as species (the modified site was not originally found in human
samples) and the change of protein sequence (the previous protein sequence is
incomplete, and the protein sequence may be prolonged with the development of protein
sequencing technology). When naming, we will use the "numbers" in historical reference
to keep the sites consistent with the reports. The antibody binds to the following
modification sequence (lowercase letters are modification sites):ASyMP

Target Information

Protein Name Leptin receptor

Organism	Gene ID	UniProt ID
Human	<u>3953;</u>	<u>P48357;</u>
Mouse	<u>16847;</u>	<u>P48356;</u>

CellularCell membrane ; Single-pass type I membrane protein . Basolateral cell membrane .;Localization[Isoform E]: Secreted .

- **Tissue specificity** Isoform A is expressed in fetal liver and in hematopoietic tissues and choroid plexus. In adults highest expression in heart, liver, small intestine, prostate and ovary. Low level in lung and kidney. Isoform B is highly expressed in hypothalamus, but also in skeletal muscle. Detected in fundic and antral epithelial cells of the gastric mucosa (PubMed:19159218). Isoform B and isoform A are expressed by NK cells (at protein level) (PubMed:12504075).
- **Function** Domain: The box 1 motif is required for JAK interaction and/or activation., Domain: The cytoplasmic domain may be essential for intracellular signal transduction by activation of JAK tyrosine kinase and STATs., Domain: The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding, Function: Receptor for obesity factor (leptin). On ligand binding, mediates signaling through IAK2/STAT3. Involved in the regulation of fat metabolism and, in a hematopoietic pathway, required for normal lymphopoiesis. May play a role in reproduction. Can also mediate the ERK/FOS signaling pathway., PTM:On ligand binding, phosphorylated on two conserved C-terminal tyrosine residues (isoform B only) by JAK2. Tyr-986 is required for complete binding and activation of PTPN11, ERK/FOS activation and, for interaction with SOCS3 (By similarity). Phosphorylation on Tyr-1141 is required for STAT3 binding/activation., similarity: Belongs to the type I cytokine receptor family. Type 2 subfamily., similarity: Contains 1 lg-like (immunoglobulin-like) domain., similarity: Contains 4 fibronectin type-III domains., subunit: On leptin stimulation, homodimerizes. The phosphorylated receptor binds a number of SH2 domain-containing proteins such as STAT3, PTPN11, and SOCS3 (By similarity). Interaction with SOCS3 inhibits LRb signaling.,tissue specificity: Isoform A is expressed in fetal liver and in hematopoietic tissues and choroid plexus. In adults highest expression in heart, liver, small intestine, prostate and ovary. Low level in lung and kidney. Isoform B is highly expressed in hypothalamus.,

Validation Data



Western Blot analysis of extracts from K562 cells, using Phospho-Ob-R (Y1141) Polyclonal Antibody.

Contact information

order.cn@immunoway.com
support.cn@immunoway.com
400-8787-807(China)
http://www.immunoway.com.cn
2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Ob-R (Phospho Tyr1141) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents