

Vimentin (Phospho Tyr38) Rabbit pAb

CatalogNo: YP1209 Orthogonal Validated 👀

Key Features

Host Species Reactivity Applications
• Rabbit • Human:Y38,Mouse:Y38,Rat:Y38 • WB,ELISA

MW Isotype
• 51kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-10000 ELISA 1:10000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human Vimentin

(Phospho-Tyr38)

Specificity This antibody detects endogenous levels of Vimentin at Human:Y38;Mouse:Y38;Rat:Y38,

It doesn't reacte with total protein. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RTySL

| Target Information

Gene name

VIM

Protein Name

vimentin

Organism	Gene ID	UniProt ID
Human	<u>7431;</u>	<u>P08670</u> ;
Mouse	22352;	<u>P20152;</u>
Rat	<u>81818;</u>	<u>P31000;</u>

Cellular Localization


Cytoplasm . Cytoplasm, cytoskeleton . Nucleus matrix . Cell membrane .

Tissue specificity Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.


Function

Function: Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells., online information: Vimentin entry, PTM: One of the most prominent phosphoproteins in various cells of mesenchymal origin. Phosphorylation is enhanced during cell division, at which time vimentin filaments are significantly reorganized..sequence Caution:Intron retention..similarity:Belongs to the intermediate filament family, subunit: Homopolymer. Interacts with HCV core protein. Interacts with LGSN and SYNM., tissue specificity: Highly expressed in fibroblasts, some expression in T- and Blymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.,


Validation Data



Western Blot analysis of HELA cells using Antibody diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Vimentin (Phospho-Ser38) Antibody



Western blot analysis of lysates from 293 cells treated with paclitaxel, using p-Vimentin (Phospho-Ser38) antibody.

| Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Vimentin (Phospho

Tyr38) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents