

## IGF-IR (Phospho Tyr1161) Rabbit pAb

CatalogNo: YP1219 **Orthogonal Validated** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA

#### MW

- pro:155kD, recetor beta:95kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-2000**

**ELISA 1:10000-20000**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized phospho derived from human IGF1R (Phospho-Tyr1161)

**Specificity** This detects endogenous levels of IGF1R (Phospho-Tyr1161). The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): DlyET

---

## | Target Information

**Gene name** IGF1R

**Protein Name** Insulin-like growth factor 1 receptor (Insulin-like growth factor I receptor) (IGF-I receptor) (CD antigen CD221) [Cleaved into: Insulin-like growth factor 1 receptor alpha chain; Insulin-like growth factor 1 receptor beta chain]


Organism	Gene ID	UniProt ID
Human	<a href="#">3480</a> ;	<a href="#">P08069</a> ;
Mouse	<a href="#">16001</a> ;	<a href="#">Q60751</a> ;
Rat	<a href="#">25718</a> ;	<a href="#">P24062</a> ;

**Cellular Localization** Cell membrane ; Single-pass type I membrane protein .

**Tissue specificity** Found as a hybrid receptor with INSR in muscle , heart , kidney , adipose tissue , skeletal muscle , hepatoma , fibroblasts , spleen and placenta (at protein level) . Expressed in a variety of tissues. Overexpressed in tumors , including melanomas , cancers of the colon , pancreas prostate and kidney.

**Function** Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Disease:Defects in IGF1R may be a cause in some cases of resistance to insulin-like growth factor 1 (IGF1 resistance) [MIM:270450]. IGF1 resistance is a growth deficiency disorder characterized by intrauterine growth retardation and poor postnatal growth accompanied with increased plasma IGF1. ,enzyme regulation:Autophosphorylation activates the kinase activity. ,Function:This receptor binds insulin-like growth factor 1 (IGF1) with a high affinity and IGF2 with a lower affinity. It has a tyrosine-protein kinase activity , which is necessary for the activation of the IGF1-stimulated downstream signaling cascade. When present in a hybrid receptor with INSR , binds IGF1. PubMed:12138094 shows that hybrid receptors composed of IGF1R and INSR isoform Long are activated with a high affinity by IGF1 , with low affinity by IGF2 and not significantly activated by insulin , and that hybrid receptors composed of IGF1R and INSR isoform Short are activated by IGF1 , IGF2 and insulin. In contrast , PubMed:16831875 shows that hybrid receptors composed of IGF1R and INSR isoform Long and hybrid receptors composed of IGF1R and INSR isoform Short have similar binding characteristics , both bind IGF1 and have a low affinity for insulin. ,online information:IGF-1 receptor entry ,PTM:Phosphorylation of Tyr-980 is required for IRS1- and SHC1-binding. ,PTM:The cytoplasmic domain of the beta subunit is autophosphorylated on tyrosine residues in response to insulin-like growth factor I (IGF I) . ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 3 fibronectin type-III domains. ,subunit:Tetramer of 2 alpha and 2 beta chains linked by disulfide bonds. The alpha chains contribute to the formation of the ligand-binding domain , while the beta chain carries the kinase domain. Interacts with PIK3R1 and with the PTB/PID domains of IRS1 and SHC1 in vitro when autophosphorylated on tyrosine residues. Forms a hybrid receptor with INSR , the hybrid is a tetramer consisting of 1 alpha chain and 1 beta chain of INSR and 1 alpha chain and 1 beta chain of IGF1R. ,tissue specificity:Found as a hybrid receptor with INSR in muscle , heart , kidney , adipose tissue , skeletal muscle , hepatoma , fibroblasts , spleen and placenta (at protein level) . Expressed in a variety of tissues. ,

## Validation Data



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody (catalog#:RS0002) was diluted at 1:20000

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**IGF-IR (Phospho Tyr1161) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)