

## RIP2 (Phospho Ser176) Rabbit pAb

CatalogNo: YP1223

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB, ELISA, IHC

#### MW

- 61kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-2000**

**IHC 1:50-300**

**ELISA 1:2000-20000**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized phospho derived from human RIPK2 (Phospho-Ser176)

**Specificity** This detects endogenous levels of RIPK2 (Phospho-Ser176). The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): SLsQS

## Target Information

**Gene name** RIPK2 CARDIAK RICK RIP2 UNQ277/PRO314/PRO34092

**Protein Name** Receptor-interacting serine/threonine-protein kinase 2 (CARD-containing interleukin-1 beta-converting enzyme-associated kinase) (CARD-containing IL-1 beta ICE-kinase) (RIP-like-interacting CLARP kinase) (Receptor-interacting protein 2) (RIP-2) (Tyrosine-protein kinase RIPK2) (EC 2.7.10.2)


Organism	Gene ID	UniProt ID
Human	<a href="#">8767</a> ;	<a href="#">O43353</a> ;
Mouse	<a href="#">192656</a> ;	<a href="#">P58801</a> ;

**Cellular Localization** Cytoplasm .


**Tissue specificity** Detected in heart , brain , placenta , lung , peripheral blood leukocytes , spleen , kidney , testis , prostate , pancreas and lymph node.

**Function** Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Activates pro-caspase-1 and pro-caspase-8. Potentiates CASP8-mediated apoptosis. Activates NF-kappa-B. ,PTM:Autophosphorylated. Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. ,similarity:Contains 1 CARD domain. ,similarity:Contains 1 protein kinase domain. ,subunit:Binds to CFLAR/CLARP and CASP1 via their CARD domains. Binds to BIRC3/c-IAP1 and BIRC2/c-IAP2 , TRAF1 , TRAF2 , TRAF5 and TRAF6. May be a component of both the TNFRSF1A and TNFRSF5/CD40 receptor complex. ,tissue specificity:Detected in heart , brain , placenta , lung , peripheral blood leukocytes , spleen , kidney , testis , prostate , pancreas and lymph node. ,

## Validation Data



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody (catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**RIP2 (Phospho Ser176) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)