

GRK 2 (Phospho Ser685) Rabbit pAb

CatalogNo: YP1225 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 80kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:10000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho derived from human GRK2 (Phospho-Ser685)

Specificity This detects endogenous levels of GRK2 (Phospho-Ser685). The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): RGSAN

Target Information

Gene name ADRBK1 BARK BARK1 GRK2

Protein Name Beta-adrenergic receptor kinase 1 (Beta-ARK-1) (G-protein coupled receptor kinase 2)


Organism	Gene ID	UniProt ID
Human	156 ;	P25098 ;
Mouse		Q99MK8 ;
Rat	25238 ;	P26817 ;

Cellular Localization Cytoplasm . Cell membrane . Cell junction , synapse , postsynapse . Cell junction , synapse , presynapse .

Tissue specificity Expressed in peripheral blood leukocytes.

Function Catalytic activity:ATP + [beta-adrenergic receptor] = ADP + [beta-adrenergic receptor] phosphate. ,Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,Function:Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors , probably inducing a desensitization of them. ,online information:Beta adrenergic receptor kinase entry ,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. GPRK subfamily. ,similarity:Contains 1 AGC-kinase C-terminal domain. ,similarity:Contains 1 PH domain. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 RGS domain. ,subunit:Interacts with GIT1 (By similarity) . Interacts with , and phosphorylates chemokine-stimulated CCR5. ,tissue specificity:Expressed in peripheral blood leukocytes. ,

Validation Data



Western blot analysis of CACO2 lysate, antibody was diluted at 1000. Secondary antibody (catalog#:RS0002) was diluted at 1:20000

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
GRK 2 (Phospho Ser685) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)