

SMC3 (Phospho Ser1083) Rabbit pAb

CatalogNo: YP1228

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, WB

MW

- 145kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:50-200

WB 1:500-2000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human Bamacan (Phospho-Ser1083)

Specificity This antibody detects endogenous phospho levels of Bamacan (Phospho-Ser1083) at Human:S1083, Mouse:S1083, Rat:S1083. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SGsQS

Target Information

Gene name SMC3 BAM BMH CSPG6 SMC3L1

Protein Name Bamacan (Phospho-Ser1083)

Organism	Gene ID	UniProt ID
Human	9126 ;	Q9UQE7 ;
Mouse		Q9CW03 ;
Rat		P97690 ;

Cellular Localization


Nucleus . Chromosome . Chromosome , centromere . Associates with chromatin. Before prophase it is scattered along chromosome arms. During prophase , most of cohesin complexes dissociate from chromatin probably because of phosphorylation by PLK , except at centromeres , where cohesin complexes remain. At anaphase , the RAD21 subunit of the cohesin complex is cleaved , leading to the dissociation of the complex from chromosomes , allowing chromosome separation. The phosphorylated form at Ser-1083 is preferentially associated with unsynapsed chromosomal regions (By similarity) . .

Tissue specificity B-cell ,Epithelium ,Eye ,Neuron ,Umbilical cord blood ,

Function

Caution:Was originally isolated as a proteoglycan protein (explaining its name) . Although not excluded , such secreted function is not clear. ,Disease:Defects in SMC3 are the cause of Cornelia de Lange syndrome type 3 (CDLS3) [MIM:610759]. CDLS is a dominantly inherited multisystem developmental disorder characterized by growth and cognitive retardation , abnormalities of the upper limbs , gastroesophageal dysfunction , cardiac , ophthalmologic and genitourinary anomalies , hirsutism , and characteristic facial features. CDSL3 is a mild form with absence of major structural anomalies typically associated with CDLS. The phenotype in some instances approaches that of apparently non-syndromic mental retardation. ,Domain:The flexible hinge domain , which separates the large intramolecular coiled coil regions , allows the heterotypic interaction with the corresponding domain of SMC1A or SMC1B , forming a V-shaped heterodimer. The two heads of the heterodimer are then connected by different ends of the cleavable RAD21 protein , forming a ring structure. ,Function:Involved in chromosome cohesion during cell cycle and in DNA repair. Central component of cohesin complex. The cohesin complex is required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At anaphase , the complex is cleaved and dissociates from chromatin , allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis and in chromosome movement. ,PTM:Phosphorylated upon DNA damage , probably by ATM or ATR. ,similarity:Belongs to the SMC family. SMC3 subfamily. ,subcellular location:Associates with chromatin. Before prophase it is scattered along chromosome arms. During prophase , most of cohesin complexes dissociate from chromatin probably because of phosphorylation by PLK , except at centromeres , where cohesin complexes remain. At anaphase , the RAD21 subunit of the cohesin complex is cleaved , leading to the dissociation of the complex from chromosomes , allowing chromosome separation. ,subunit:Interacts with MXI1 , MXD3 and MXD4. Interacts with SYCP2. Found in a complex with SMC1A , CDCA5 and RAD21 , PDS5A/APRIN and PDS5B/SCC-112 (By similarity) . Forms a heterodimer with SMC1A or SMC1B in cohesin complexes. Cohesin complexes are composed of the SMC1 (SMC1A or SMC1B) and SMC3 heterodimer attached via their hinge domain , RAD21 which link them , and one STAG protein (STAG1 , STAG2 or STAG3) , which interacts with RAD21. Also found in meiosis-specific cohesin complexes. Interacts with NUMA1 , and forms a ternary complex with KIF3B and KIFAP3 , suggesting a function in tethering the chromosomes to the spindle pole and in chromosome movement. ,

Validation Data



Immunohistochemical analysis of paraffin-embedded human brain tumor. 1, Antibody was diluted at 1:200 (4°C overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SMC3 (Phospho Ser1083) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)