

Trk B (Phospho Y817) Rabbit pAb

CatalogNo: YP1247 **Orthogonal Validated** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- IHC, IF, WB

MW

- 145kd or 92kd (with limited samples) (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:50-200

WB 1:500-2000

IF 1:50-200

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human Trk B (Phospho-Y817)

Specificity This antibody detects endogenous phospho levels of Trk B (Phospho-Y817) Polyclonal Antibody at Human, Mouse, Rat. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PVyLD

Target Information

Gene name NTRK2 TRKB

Protein Name Trk B (Phospho-Y817) Polyclonal Antibody

Organism	Gene ID	UniProt ID
Human	4915;	Q16620;

Cellular Localization

Cell membrane ; Single-pass type I membrane protein . Endosome membrane ; Single-pass type I membrane protein . Early endosome membrane . Cell projection , axon . Cell projection , dendrite . Cytoplasm , perinuclear region . Cell junction , synapse , postsynaptic density . Internalized to endosomes upon ligand-binding. .


Tissue specificity

Isoform TrkB is expressed in the central and peripheral nervous system. In the central nervous system (CNS) , expression is observed in the cerebral cortex , hippocampus , thalamus , choroid plexus , granular layer of the cerebellum , brain stem , and spinal cord. In the peripheral nervous system , it is expressed in many cranial ganglia , the ophthalmic nerve , the vestibular system , multiple facial structures , the submaxillary glands , and dorsal root ganglia. Isoform TrkB-T1 is mainly expressed in the brain but also detected in other tissues including pancreas , kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in the brain.

Function

Alternative products:Additional isoforms seem to exist ,Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,Function:Receptor for brain-derived neurotrophic factor (BDNF) , neurotrophin-3 and neurotrophin-4/5 but not nerve growth factor (NGF) . Involved in the development and/or maintenance of the nervous system. This is a tyrosine-protein kinase receptor. Known substrates for the TRK receptors are SHC1 , PI-3 kinase , and PLC-gamma-1. ,PTM:Ligand-mediated auto-phosphorylation. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains. ,similarity:Contains 2 LRR (leucine-rich) repeats. ,subunit:Exists in a dynamic equilibrium between monomeric (low affinity) and dimeric (high affinity) structures. Binds SH2B2. Interacts with SQSTM1 and KIDINS220. ,tissue specificity:Isoform TrkB is widely expressed , mainly in the nervous tissue. In the CNS , expression is observed in the cerebral cortex , hippocampus , thalamus , choroid plexus , granular layer of the cerebellum , brain stem , and spinal cord. In the peripheral nervous system , it is expressed in many cranial ganglia , the ophtalmic nerve , the vestibular system , multiple facial structures , the submaxillary glands , and dorsal root ganglia. Isoform TrkB-T1 is expressed in multiple tissues , mainly in brain , pancreas , kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in brain. ,

Validation Data



The SLITRK4-CNPY3 axis promotes liver metastasis of gastric cancer by enhancing the endocytosis and recycling of TrkB in tumour cells CELLULAR ONCOLOGY Li Jun WB Human BGC823 cell,MKN28 cell

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**Trk B (Phospho
Y817) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)