

## FoxO4 (Phospho Ser193) Rabbit pAb

CatalogNo: YP1343

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 55kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

WB 1:1000-2000

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized phospho peptide around human FoxO4 (Ser193)

**Specificity** This antibody detects endogenous levels of Human FoxO4 (phospho-Ser193)

### Target Information

**Gene name** FOXO4 AFX AFX1 MLLT7

**Protein Name** Forkhead box protein O4 (Ser193)

Organism	Gene ID	UniProt ID
Human	<a href="#">4303</a> ;	<a href="#">P98177</a> ;
Mouse	<a href="#">54601</a> ;	<a href="#">Q9WVH3</a> ;

**Cellular Localization** Cytoplasm. Nucleus. When phosphorylated , translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation. Monoubiquitination increases nuclear localization. When deubiquitinated , translocated from nucleus to cytoplasm.

**Tissue specificity** Heart , brain , placenta , lung , liver , skeletal muscle , kidney and pancreas. Isoform zeta is most abundant in the liver , kidney , and pancreas.

**Function** Disease:A chromosomal aberration involving FOXO4 is found in acute leukemias. Translocation t (X;11) (q13;q23) with MLL/HRX. The result is a rogue activator protein. ,Function:Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGF1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle. ,pharmaceutical:A constitutively active FOXO4 mutant where phosphorylation sites Thr-32 , Ser-187 and Ser-262 have been mutated to alanine may have therapeutic potential in ERBB2/HER2-overexpressing cancers as it inhibits ERBB2-mediated cell survival , transformation and tumorigenicity. ,PTM:Acetylation by CBP , which is induced by peroxidase stress , inhibits transcriptional activity. Deacetylation by SIRT1 is NAD-dependent and stimulates transcriptional activity. ,PTM:Phosphorylation by PKB/AKT1 inhibits transcriptional activity and is responsible for cytoplasmic localization. ,similarity:Contains 1 fork-head DNA-binding domain. ,subcellular location:When phosphorylated , translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation. ,subunit:Interacts with CBP , MYOCD , SIRT1 , SRF and YWHAZ. Acetylated by CBP and deacetylated by SIRT1. Binding of YWHAZ inhibits DNA-binding. ,tissue specificity:Heart , brain , placenta , lung , liver , skeletal muscle , kidney and pancreas. Isoform zeta is most abundant in the liver , kidney , and pancreas. ,

## Validation Data

## Contact information

Orders: [order.cn@immunoway.com](mailto:order.cn@immunoway.com)  
Support: [support.cn@immunoway.com](mailto:support.cn@immunoway.com)  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**FoxO4 (Phospho Ser193) Rabbit pAb**