

## PAK1/2 (Phospho Ser199/204) Rabbit pAb

CatalogNo: YP1431

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse, Rat

#### Applications

- WB

#### MW

- 60kD (Observed)

#### Isotype

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

WB 1:1000-2000

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized phospho peptide around human PAK1 (Ser199 and 204)

**Specificity** This antibody detects endogenous levels of PAK1/2 only when phosphorylated at Ser199/204. This antibody does not recognize phosphorylated at other sites.

### Target Information

**Protein Name** PAK1/2 (Ser199/204)

Organism	Gene ID	UniProt ID
Human	<a href="#">5058</a> ;	<a href="#">Q13153</a> ;
Mouse		<a href="#">O88643</a> ;
Rat	<a href="#">29431</a> ;	<a href="#">P35465</a> ;

**Cellular Localization**

Cytoplasm . Cell junction , focal adhesion . Cell projection , lamellipodium . Cell membrane . Cell projection , ruffle membrane . Cell projection , invadopodium . Nucleus , nucleoplasm . Chromosome . Cytoplasm , cytoskeleton , microtubule organizing center , centrosome . Colocalizes with RUFY3 , F-actin and other core migration components in invadopodia at the cell periphery (PubMed:25766321) . Recruited to the cell membrane by interaction with CDC42 and RAC1. Recruited to focal adhesions upon activation. Colocalized with CIB1 within membrane ruffles during cell spreading upon readhesion to fibronectin. Upon DNA damage , translocates to the nucleoplasm when phosphorylated at Thr-212 where is co-recruited with MORC2 on damaged chromatin (PubMed:23260667) . Localization to the centrosome does not depend upon the presence of gamma-tubulin (PubMed:27012601) . Localization of the active , but not inactive , protein to the adhesions and edge of lamellipodia is mediated by interaction with GIT1 (PubMed:11896197) . .

**Tissue specificity** Overexpressed in gastric cancer cells and tissues (at protein level) (PubMed:25766321) .

**Function**

Catalytic activity:ATP + a protein = ADP + a phosphoprotein. ,cofactor:Magnesium. ,enzyme regulation:Activated by binding small G proteins. Binding of GTP-bound CDC42 or RAC1 to the autoregulatory region releases monomers from the autoinhibited dimer , enables phosphorylation of Thr-423 and allows the kinase domain to adopt an active structure. Also activated by binding to GTP-bound CDC42 , independent of the phosphorylation state of Thr-423. Phosphorylation of Thr-84 by OXSR1 inhibits this activation. ,Function:The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis , potentially due to binding of CDC2L1 and CDC2L2. ,PTM:Autophosphorylated when activated by CDC42/p21 and RAC1. ,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily. ,similarity:Contains 1 CRIB domain. ,similarity:Contains 1 protein kinase domain. ,subcellular location:Recruited to focal adhesions upon activation. ,subunit:Homodimer in its autoinhibited state. Active as monomer. Interacts tightly with GTP-bound but not GDP-bound CDC42/P21 and RAC1. Binds to the caspase-cleaved p110 isoform of CDC2L1 and CDC2L2 , p110C , but not the full-length proteins. Component of cytoplasmic complexes , which also contain PXN , ARHGEF6 and GIT1. Interacts with ARHGEF7. Also interacts with CRIPAK. Interacts with NISCH. ,

---

## | Validation Data

## | Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**PAK1/2 (Phospho Ser199/204) Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)